

DESCRIPTION

Species Reactivity	Mouse/Rat
Specificity	Detects mouse and rat in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant mouse Integrin β 5 Gly24-Asn719 Accession # NP_001139356
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

CyTOF-ready	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Flow Cytometry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The β 5 integrin is a 95-110 kDa member of the integrin beta chain family, integrin superfamily of molecules. It is expressed by select cell types, including microglia, keratinocytes, fibroblasts, endothelial cells, adhesive monocytes and macrophages, retinal pigment epithelium, immature dendritic cells and colonic epithelium. The Integrin β 5 subunit forms a noncovalent heterodimer with the α v integrin subunit, and together they bind vitronectin, and the Thy-1 antigen. α v β 5 is reported to participate in phagocytosis, and act in concert with CD81, CD36 and stabilin-2 on the cell surface. When α v β 5 binds Thy-1, the heterodimer apparently interferes with cell membrane activation of TGF- β . Intracellularly, it appears to regulate actin skeleton remodeling via focal adhesion complex formation, and may contribute to cryptic pathways associated with antigen presentation. Mature mouse Integrin β 5 is a 775 amino acid (aa) type I transmembrane glycoprotein that contains a 696 aa extracellular domain (ECD) (aa 24-719) and a 56 aa cytoplasmic region. The ECD contains a 243 aa vWF-A domain (aa 136-378) and a Cys-rich tandem repeat region (aa 465-630). There is one splice variant that shows a 57 aa substitution for aa 760-798. This splice form does not appear to be expressed by the human Integrin β 5 gene. Over aa 24-719, mouse Integrin β 5 shares 91% and 97% aa sequence identity with human and rat Integrin β 5, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.