

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects endogenous human, mouse and rat BAK in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human BAK Pro20-Asn124 Accession # Q16611
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

BAK (Bcl-2 homologous antagonist/killer; also BAK1) is a 25-30 kDa member of the BCL-2 family of proteins. It is widely expressed, and participates in the apoptotic cycle. BAK is an outer mitochondrial membrane protein that is inactive as a Zn-dependent homodimer. Upon activation by p53 or tBID, BAK oligomerizes, creating a pore in the mitochondrial membrane and allowing for cytochrome C release. Human BAK is 211 amino acids (aa) in length and contains three BCL-2 homology domains (aa 74-88, 117 -136 and 169-184), a Zn-binding region (aa 160-166) and a C-terminal transmembrane segment (aa 188-205). Amino acids 67-94 mediate oligomerization of BAK. There are two potential isoform variants; one shows an alternate start site at Met96, while a second shows a deletion of aa 46-66. Over amino acids 20-124, human BAK shares 76% aa identity with mouse BAK.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.