

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Thymidine Kinase 1 Protein in direct ELISAs.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived human Thymidine Kinase 1 protein Ser2-Asn234 Accession # P04183
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide

*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Thymidine kinases are 2'-deoxythymidine kinases that phosphorylate deoxythymidine and generate deoxythymidine 5'-phosphate (1). Thymidine kinases have a key function in the synthesis of DNA and thereby in cell division, as they are part of the unique reaction chain to introduce deoxythymidine into the DNA. Two forms of thymidine kinase are present in mammalian cells, TK1 and TK2. TK1 is cell cycle-dependent and is present in the cytoplasm only in anticipation of cell division (2, 3); whereas TK2 is located in mitochondria and is cell cycle-independent (4). TK1 is synthesized by the cell during the S phase of cell division and is degraded after cell division is completed (5). TK1 normally occurs in tissue as a dimer and can be converted to more active tetrameric form by ATP binding (6). TK1 is feedback inhibited by thymidine triphosphate, the product of the further phosphorylation of thymidine (7). Because tumor cells replicate much more frequently than normal cells and requires faster DNA synthesis and higher TK1 activity, TK1 is a cancer marker especially for hematologic malignancies (8, 9). In clinical chemistry TK1 is used as a proliferation marker in the diagnosis, control of treatment and follow-up of malignant disease (10). In addition, thymidine kinase is required for the action of many antiviral drugs (11). The enzymatic activity of recombinant human TK1 is measured using a phosphatase-coupled method (12).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.