

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human MUC-4 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human MUC-4 Asn1137-Glu1317 Accession # Q99102
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

MUC-4 (Mucin-4), also called ASGP (Ascites Sialoglycoprotein) is a highly glycosylated type I transmembrane glycoprotein that may be up to 950 kDa in its full-length, fully glycosylated form. Human MUC-4 cDNA encodes 2169 amino acids (aa) with a 28 aa signal sequence and a cleavage site that creates a 1416 aa soluble, extracellular alpha chain and a 725 aa single-pass transmembrane beta chain. Between aa 1072-1317 within the alpha chain, human MUC-4 shares 69% aa sequence identity with mouse and rat MUC-4. At least 14 soluble or transmembrane splice variants of 1102-2117 aa have been described, 5 of which contain the full sequence used as an immunogen. MUC-4 can serve as a ligand for the oncogenic receptor ErbB2 and a modulator of its phosphorylation and signaling. MUC-4 is frequently aberrantly expressed in epithelial tumors and can promote tumor growth and metastasis.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.