

## DESCRIPTION



<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human GM130/GOLGA2 in direct ELISAs and Western blots.
<b>Source</b>	Polyclonal Sheep IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human GM130/GOLGA2 Met528-Gln606 Accession # Q08379
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	0.25 µg/mL	See Below
<b>Immunocytochemistry</b>	5-15 µg/mL	See Below

## DATA

<p><b>Western Blot</b></p>  <p><b>Detection of Human GM130/GOLGA2 by Western Blot.</b> Western blot shows lysates of MCF-7 human breast cancer cell line and ZR-75 human breast cancer cell line. PVDF membrane was probed with 0.25 µg/mL of Sheep Anti-Human GM130/GOLGA2 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF8199) followed by HRP-conjugated Anti-Sheep IgG Secondary Antibody (Catalog # HAF016). A specific band was detected for GM130/GOLGA2 at approximately 130 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.</p>	<p><b>Immunocytochemistry</b></p>  <p><b>GM130/GOLGA2 in HeLa Human Cell Line.</b> GM130/GOLGA2 was detected in immersion fixed HeLa human cervical epithelial carcinoma cell line using Sheep Anti-Human GM130/GOLGA2 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF8199) at 1.7 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Sheep IgG Secondary Antibody (red; Catalog # NL010) and counterstained with DAPI (blue). Specific staining was localized to Golgi apparatus vesicles. View our protocol for <a href="#">Fluorescent ICC Staining of Cells on Coverslips</a>.</p>
---	--

## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

GM130, also known as Golgin subfamily A member 2 and GOLGA2, is a 130 kDa member of the Golgin protein family. GM130/GOLGA2 is a component of the Golgi membrane involved in maintaining cis-Golgi structure. GM130/GOLGA2 protein is thought to play roles in the stacking of Golgi cisternae and in vesicular transport. Several alternatively spliced transcript variants of this gene have been predicted. Human GM130/GOLGA2 protein is 1002 amino acids (aa) in length, and over aa 528-606, shares 90% sequence identity with its mouse and rat orthologs.