

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse Caspase-7 precursor and the small Caspase-7 subunit that is generated during proteolytic activation.
Source	Polyclonal Rabbit IgG
Purification	Antigen Affinity-purified
Immunogen	KLH-coupled mouse Caspase-7 synthetic peptide RHFESQSDDPRFNEKC
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Knockout Validated	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Caspase-7 (Cysteine-aspartic acid protease 7/Casp7; also CMH-1, ICE-LAP3 and Mch3) is a 32 kDa member of the peptidase C14A/IL-1 β -converting family of enzymes (1, 2, 3). It is widely expressed, except in brain, and is best known as an integral component of the apoptotic cascade. Caspase-7 is considered to be an executioner caspase, as a downstream mediator of apoptotic-associated proteolysis (2, 3). Upon activation, Caspase-7 is known to utilize a Cys residue to cleave multiple substrates, including PARP, procaspase 6, Gas2 and calpstatin (1). Human procaspase-7 is a 34-36 kDa, 303 amino acid (aa) protein (4, 5, 6). Normally, it is an inactive homodimer (1, 2, 7, 8). But following an upstream signal that activates processing proteases, procaspase-7 undergoes proteolytic cleavage to generate an N-terminal 23 aa propeptide, a 175 aa p20/20 kDa subunit (aa 24-198), and a 105 aa C-terminal p12/12 kDa subunit (5). The p20 and p12 subunits noncovalently heterodimerize, and subsequently associate with another p20/p12 heterodimer to form an active antiparallel homodimer. Additional processing of p20 may remove aa 24-36 to generate p18, while additional processing of p12 will remove aa 199-206 to generate p11 (9, 10). Multiple proteases can use Caspase-7 as a substrate, and include caspase-1, -3, -8, and -10, granzyme B, calpain-1 and Caspase-7 itself (3, 6, 9, 11). Caspase-7 is found in both cytosol and nucleus, and possesses a potential KKKK nuclear localization signal between aa 38-41 that likely undergoes sumoylation (9, 12). There are two potential isoform variants, one which shows an alternate start site 33 aa upstream of the standard start site, and a second that shows a 105 aa substitution for aa 149-303. Human and mouse Caspase-7 are 82% aa identical at the amino acid level.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.