

DESCRIPTION




Species Reactivity	Mouse
Specificity	Detects mouse Pref-1/DLK1/FA1 in direct ELISAs and Western blots. In direct ELISAs, approximately 75% cross-reactivity with recombinant human Pref-1 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Pref-1/DLK1/FA1 Ala24-Gln305 Accession # Q09163
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
Immunohistochemistry	5-15 µg/mL	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Pref-1 (Preadipocyte factor 1, Protein delta homolog 1, DLK1, FA1 and Fetal antigen 1) is a 45-60 kDa transmembrane glycoprotein that is highly expressed in fetal liver, placenta, adult adrenal gland, brain, testis and ovary. Expression of Pref-1 is elevated in liver after birth but starts to decline around postnatal day 16. It contains 6 EGF-like domains and is involved in embryonic skeletal system development. Pref-1 inhibits preadipocyte proliferation by regulating their entry into G1/S-phase and represses preadipocyte differentiation. It is a master regulator of preadipocyte homeostasis and adipose tissue expansion. Pref-1 manipulation may, therefore, be utilized in obesity treatments. Mouse Pref-1 shares 81% aa identity with human Pref-1.