

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human NAIP expressed in Sf21 insect cells.
Source	Polyclonal Rabbit IgG
Purification	Antigen Affinity-purified
Immunogen	KLH-coupled human NAIP synthetic peptide YLSLSSTRPDEGLASIIC Accession # NP_004527
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Neuronal apoptosis inhibitory protein (NAIP; also Baculoviral IAP repeat-containing protein 1) is a 160 kDa member of the inhibitor of apoptosis family of proteins (also known as BIRC proteins). Human NAIP is 1403 amino acids (aa) in length. It contains three distinct regions: an N-terminal cluster of three baculoviral inhibitory repeat (BIR) domains, a central nucleotide binding oligomerization domain (NOD), and a C-terminal leucine-rich repeat (LRR) domain. Human NAIP shares 68% aa identity with mouse NAIP. NAIP is expressed in motor neurons, but not in sensory neurons. It is also expressed in the liver, placenta and to a lesser extent in the spinal cord. NAIP prevents motor neuron apoptosis, and defects in NAIP have been found in individuals with spinal muscular atrophy.