

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human and mouse Caspase-10 (FLICE 2) precursor in Western blots. Sensitivity for detection of the small subunit is not known.
Source	Polyclonal Rabbit IgG
Purification	Antigen Affinity-purified
Immunogen	KLH-coupled human Caspase-10 (FLICE 2) synthetic peptide KRTVWGAKQISATSLPTAC
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Caspase-10 (Cysteine-aspartic acid protease 10, FLICE 2 and Mch4) is a member of the peptidase C14A family of enzymes. It is widely expressed, being found in blood and epithelial cells. Caspase-10 appears to be an initiator caspase that acts on Pro-Caspases-3, -4, -6, -7, and -9. Human Pro-Caspase-10a is a 59 kDa, 521 amino acid (aa) protein. It contains two N-terminal death domains. Cleavage generates p43 plus the small mature p12 subunit. Cleavage of p43 generates a prodomain p25 and large mature p17 subunit. P17 and p12 noncovalently associate to form a 29 kDa mature heterodimer. There are at least four isoform variants. There is no apparent mouse Casp-10 gene.