

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-12 Rβ1 in direct ELISAs and Western blots. In these formats, less than 2% cross-reactivity with recombinant mouse IL-12 Rβ1 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human IL-12 Rβ1 Cys24-Glu540 Accession # P42701
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Interleukin 12 (IL-12) is a key mediator of cellular-immunity and induces the differentiation of Th1 cells from precursor T helper cells. The biological activities of IL-12 are mediated through the high-affinity receptor complex composed of two subunits designated IL-12 Rβ1 and IL-12 Rβ2. Individually, IL-12 Rβ1 and IL-12 Rβ2 bind IL-12 with low affinity. Co-expression of both subunits confers high-affinity binding and is required for IL-12 activity. Both IL-12 receptor subunits are type I membrane proteins that share similarities with the gp130/G-CSF R subgroup in the cytokine receptor superfamily. IL-12 Rβ1 cDNA encodes a 662 amino acid (aa) residue protein with a putative 23 aa residue signal peptide that is cleaved to generate the mature protein with a 522 aa residue extracellular domain, a 25 aa residue transmembrane domain and a 92 aa residue cytoplasmic region. Expression of IL-12 Rβ1 has been detected in activated T cells, NK cells and B cells. The expression of IL-12 Rβ2 is more restricted and appears to be limited to Th2 cells.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.