

DESCRIPTION



Species Reactivity	Human/Mouse/Rat
Specificity	Detects human, mouse, and rat RSK1 when phosphorylated at S221, and human and mouse RSK2 when phosphorylated at S227.
Source	Polyclonal Rabbit IgG
Purification	Antigen Affinity-purified
Immunogen	Phosphopeptide containing human RSK1 S221 site
Formulation	Supplied as a 0.2 µm filtered solution in PBS with Sodium Azide. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.5 µg/mL	See Below
Immunohistochemistry	5-15 µg/mL	See Below

DATA

Western Blot	Immunohistochemistry
 <p>Detection of Human Phospho-RSK1 (S221) and RSK2 (S227) by Western Blot. Western blot shows lysates of HeLa human cervical epithelial carcinoma cell line untreated (-) or treated (+) with 200 nM PMA for for the indicated times. PVDF membrane was probed with 0.5 µg/mL of Rabbit Anti-Human/Mouse/Rat Phospho-RSK1 (S221)/RSK2 (S227) Antigen Affinity-purified Polyclonal Antibody (Catalog # AF892), followed by HRP-conjugated Anti-Rabbit IgG Secondary Antibody (Catalog # HAF008). A specific band was detected for Phospho-RSK1 (S221) and RSK2 (S227) at approximately 90 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.</p>	 <p>RSK1/RSK2 in Mouse Brain. RSK1/RSK2 was detected in perfusion fixed frozen sections of mouse brain (Purkinje cells in cerebellum) using 15 µg/mL Rabbit Anti-Human/Mouse/Rat Phospho-RSK1/RSK2 (RSK1 S221, RSK2 S227) Antigen Affinity-purified Polyclonal Antibody (Catalog # AF892) overnight at 4 °C. Tissue was stained with the Anti-Rabbit HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS005) and counterstained with hematoxylin (blue). View our protocol for Chromogenic IHC Staining of Frozen Tissue Sections.</p>

PREPARATION AND STORAGE

Shipping	The product is shipped with dry ice or equivalent. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C, as supplied. ● 1 month, 2 to 8 °C under sterile conditions after opening. ● 6 months, -20 to -70 °C under sterile conditions after opening.