

## DESCRIPTION


<b>Species Reactivity</b>	Human/Mouse
<b>Specificity</b>	Detects mouse ADAM9 Ectodomain in direct ELISAs and Western blots. In Western blots no cross-reactivity with the Ectodomain of recombinant mouse ADAM10 and rhADAM8, 15, and 17 (TACE) is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant mouse ADAM9 Ala206-Asp697 Accession # Q61072
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	1 µg/mL	See Below
<b>Flow Cytometry</b>	0.25 µg/10 <sup>6</sup> cells	Mouse splenocytes
<b>Immunohistochemistry</b>	5-15 µg/mL	Perfusion fixed frozen sections of mouse testis
<b>Immunoprecipitation</b>	25 µg/mL	Conditioned cell culture medium spiked with Recombinant Mouse ADAM9 (Catalog # 949-AD), see our available <a href="#">Western blot detection antibodies</a>
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<p><b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b></p> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

ADAM9, also known as MDC9 or meltrin  $\gamma$ , is a member of the ADAM family that contains a disintegrin and metalloprotease-like domain (1). Like other membrane-anchored ADAMs, ADAM9 consists of a pro domain with a cysteine switch and furin cleavage sequence, a catalytic domain with the zinc-binding site and Met-turn expected for reprolysins, a disintegrin-like domain, a cysteine-rich domain, an EGF-like domain, a transmembrane domain, and the cytoplasmic domain. ADAM9 is able to cleave peptides corresponding to cleavage sites of tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), the p75-TNF receptor, the  $\beta$ -amyloid protein precursor, and the c-kit ligand-1, implying that it may participate in shedding of these membrane proteins (2). In fact, ADAM9 has been shown to shed membrane-anchored heparin-binding EGF-like growth factor (3). In addition, it also cleaves oxidized insulin B-chain and fibronectin (2, 4). Besides its catalytic activity, ADAM9 functions as an adhesion molecule through binding of its disintegrin domain to integrins such as  $\alpha_v\beta_5$  and  $\alpha_6\beta_1$  (5, 6). The cytoplasmic domain of ADAM9 interacts with Src homology 3 (SH3)-containing proteins and protein kinase C, and may mediate different signaling pathways (3, 7). ADAM9 is widely expressed in tissues (8).

## References:

1. Moss, M.L. *et al.* (2001) *Drug Discov. Today* **6**:417.
2. Roghani, M. *et al.* (1999) *J. Biol. Chem.* **274**:3531.
3. Izumi, Y. *et al.* (1998) *EMBO J.* **17**:7260.
4. Schwettmann, L. and H. Tschesche (2001) *Protein. Expr. Purif.* **21**:65.
5. Nath, D. *et al.* (2000) *J. Cell Sci.* **113**:2319.
6. Zhou, M. *et al.* (2001) *Biochem. Biophys. Res. Comm.* **280**:574.
7. Howard, L. *et al.* (1999) *J. Biol. Chem.* **274**:31693.
8. Weskamp, G. *et al.* (1996) *J. Cell Biol.* **132**:717.