

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Angiopoietin-4 in direct ELISAs and Western blots. In direct ELISAs, less than 5% cross-reactivity with recombinant human (rh) ANG-1, rhANG-2 and recombinant mouse ANG-3 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Angiopoietin-4 Gln23-Ile503 Accession # Q9Y264
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Blockade of Receptor-ligand Interaction	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

GITR (glucocorticoid-induced TNF receptor superfamily-related protein, also named AITR, activation-inducible TNF receptor superfamily-related protein) and GITR ligand (GITRL) are novel members of the TNF receptor (TNFR) and TNF superfamilies (SF) that have been designated TNFRSF18 and TNFSF18, respectively. Human GITRL cDNA encodes a 177 amino acid residues type II membrane protein. The carboxy-terminal extracellular domain shows sequence identity to TNF/TNFSF2 (21%), Fas ligand/TNFSF6 (21%), TRAIL/TNFSF10 (18%), and lymphotoxin α/TNFSF1 (18%). GITRL is constitutively expressed in human umbilical vein endothelial cells but is not expressed in resting or stimulated T cell lines, B cell lines or peripheral blood mononuclear cells. GITR, the receptor for GITRL, is expressed at low levels in peripheral blood T cells, bone marrow, thymus, spleen and lymph nodes. In contrast to mouse GITR, expression of human GITR is not induced by treatment with dexamethasone, but is up-regulated by antigen-receptor stimulation or by treatment with soluble anti-CD3 plus anti-CD28 or PMA plus ionomycin. Ligation of GITR has been found to induce nuclear factor (NF)-κB activation via TNF receptor-associated factor 2 and protect cells from TCR activation-induced cell death. It has been proposed that GITRL and GITR may modulate T lymphocyte functions in peripheral tissues.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.