

DESCRIPTION

Species Reactivity	Rat
Specificity	Detects E-Selectin/CD62E in direct ELISAs and Western blots. In direct ELISAs, approximately 15% cross-reactivity with recombinant mouse (rm) E-Selectin is observed, approximately 5% cross-reactivity with recombinant human (rh) E-Selectin.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant rat E-Selectin/CD62E Trp22-Pro494 Accession # P98105
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Adhesion Blockade	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

E-Selectin (Endothelial Leukocyte Adhesion Molecule-1, ELAM-1, CD62E), a member of the Selectin family, is a 107-115 kDa cell surface glycoprotein. It is transiently expressed on vascular endothelial cells in response to IL-1 β and TNF- α , and demonstrates peak expression at 4 hours, and decay at 24 hours, in response to activation. E-Selectin ligands, expressed on neutrophils, monocytes, and a subset of memory T cells, are sialylated, fucosylated molecules which bind to the lectin domain of E-Selectin. Immunocytochemical techniques have demonstrated the expression of E-Selectin on healthy and diseased tissue. The human and rat proteins share approximately 67% amino acid identity. The mouse and rat proteins share approximately 78% amino acid identity.

E-Selectin mediates the attachment of flowing leukocytes to the blood vessel wall during inflammation by binding to E-Selectin ligands on leukocytes. These interactions are labile and permit leukocytes to roll along the vascular endothelium in the direction of blood flow. This initial interaction is followed by a stronger interaction involving ICAM-1 and VCAM-1 that leads eventually to extravasation of the white blood cell through the blood vessel wall into the extracellular matrix tissue.

ELISA techniques have shown that detectable levels of soluble E-Selectin are present in the biological fluids of apparently normal individuals. Furthermore, a number of studies have reported that levels of E-Selectin may be elevated in subjects with a variety of pathological conditions.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.