

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse TIMP-1 in direct ELISAs and Western blots. In direct ELISAs, approximately 15% cross-reactivity with recombinant rat TIMP-1 is observed, less than 5% cross-reactivity with recombinant human (rh) TIMP-1 is observed, and less than 1% cross-reactivity with recombinant human TIMP-2 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse TIMP-1 Cys25-Arg205 Accession # P12032
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunoprecipitation	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Tissue inhibitors of metalloproteinases or TIMPs are a family of homologous proteins that regulate the activity of matrix metalloproteinases (MMPs) (1, 2). There are four known members of the family, TIMP-1, TIMP-2, TIMP-3, and TIMP-4 that have been found to exhibit multiple functions, including inhibition of active MMPs, pro-MMP activation, cell growth promotion, matrix binding, inhibition of angiogenesis and the induction of apoptosis. Structurally, TIMPs have two domains, an N-terminal domain and a C-terminal domain. Each domain consists of three disulfide-bonded loops. TIMP-1 is a glycoprotein produced by a wide range of cell types. Through its N-terminal domain, TIMP-1 inhibits active MMPs by forming a non-covalent binary complex with the MMP active site. The C-terminal domain of TIMP-1 interacts with the C-terminal domain of pro-MMP-9, which may play a role in regulating pro-MMP-9 activation.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.