

DESCRIPTION

Source *E. coli*-derived human G-CSF protein
Thr31-Pro204, with an N-terminal Met
Accession # NP_757373
Produced using non-animal reagents in an animal-free laboratory.

N-terminal Sequence Analysis Met

Predicted Molecular Mass 18.8 kDa

SPECIFICATIONS

Activity Measured in a cell proliferation assay using NFS-60 mouse myelogenous leukemia lymphoblast cells. Shirafuji, *N.et al.* (1989) Exp. Hematol. 17:116.
The ED₅₀ for this effect is 10-60 pg/mL.

Endotoxin Level <0.01 EU per 1 µg of the protein by the LAL method.

Purity >97%, by SDS-PAGE under reducing conditions and visualized by silver stain.

Formulation Lyophilized from a 0.2 µm filtered solution in Acetic Acid with Trehalose. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution Reconstitute at 0.2 mg/mL in sterile PBS.

Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage **Use a manual defrost freezer and avoid repeated freeze-thaw cycles.**

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 3 months, 2 to 8 °C under sterile conditions after reconstitution.

BACKGROUND

G-CSF is a pleiotropic cytokine best known for its specific effects on the proliferation, differentiation, and activation of hematopoietic cells of the neutrophilic granulocyte lineage. It is produced mainly by monocytes and macrophages upon activation by endotoxin, TNF-α and IFN-γ. Other cell types including fibroblasts, endothelial cells, astrocytes and bone marrow stromal cells can also secrete G-CSF after LPS, IL-1 or TNF-α activation. In addition, various carcinoma cell lines and myeloblastic leukemia cells can express G-CSF constitutively.

In humans, two distinct cDNA clones for G-CSF, encoding 207 and 204 amino acid precursor proteins, have been isolated. Both proteins have a 30 amino acid signal peptide and have identical amino acid sequences except for a three amino acid insertion (deletion) at the 35th amino acid residue from the N-terminus of the mature protein. Human G-CSF is 73% identical at the amino acid level to murine G-CSF and the two proteins show species cross-reactivity.

In vitro, G-CSF stimulates growth, differentiation and functions of cells from the neutrophil lineage. It also has blast cell growth factor activity and can synergize with IL-3 to shorten the G₀ period of early hematopoietic progenitors. Consistent with its *in vitro* functions, G-CSF has been found to play important roles in defense against infection, in inflammation and repair, and in the maintenance of steady state hematopoiesis.

MANUFACTURING SPECIFICATIONS

Animal-Free Manufacturing Conditions

Our dedicated controlled-access animal-free laboratories ensure that at no point in production are the products exposed to potential contamination by animal components or byproducts. Every stage of manufacturing is conducted in compliance with R&D Systems' stringent Standard Operating Procedures (SOPs). Production and purification procedures use equipment and media that are confirmed animal-free.

Production

- All molecular biology procedures use animal-free media and dedicated labware.
- Dedicated fermentors are utilized in committed animal-free areas.

Purification

- Protein purification columns are animal-free.
- Bulk proteins are filtered using animal-free filters.
- Purified proteins are stored in animal-free containers in a dedicated cold storage room.

Quality Assurance

- Low Endotoxin Level.
- No impairment of biological activity.
- High quality product obtained under stringent conditions.
- For *ex vivo* research or bioproduction, [additional documentation](#) can be provided.

[Please read our complete Animal-Free Statement](#)