

DESCRIPTION

Source *E. coli*-derived human IL-21 protein
Gln32-Ser162, with a N-terminal Met
Accession # Q9HBE4.3
Produced in an animal-free laboratory.

N-terminal Sequence Analysis Met

Predicted Molecular Mass 15 kDa

SPECIFICATIONS

SDS-PAGE 16 kDa, reducing conditions

Activity Measured by its ability to enhance IFN-γ secretion in NK-92 human natural killer lymphoma cells.
The ED₅₀ for this effect is ≤8 ng/mL.

Endotoxin Level <0.10 EU per 1 μg of the protein by the LAL method.

Purity >95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Formulation Lyophilized from a 0.2 μm filtered solution in PBS. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution Reconstitute at 100 μg/mL in PBS.


Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.


DATA

Bioactivity



Animal-Free™ Recombinant Human IL-21 (Catalog # AFL8879) enhances IFN-gamma secretion in NK-92 human natural killer lymphoma cells. The ED₅₀ for this effect is ≤8 ng/mL.

SDS-PAGE



2 μg/lane of Animal-Free™ Recombinant Human IL-21 (Catalog # AFL8879) was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining, showing R band at 16.4 kDa and NR band at 14.8 kDa, respectively.

BACKGROUND

Interleukin-21 (IL-21) is an approximately 14 kDa four-helix-bundle member of the family of cytokines that utilize the common gamma chain (γ_c) as a receptor subunit. γ_c is also a subunit of the receptors for IL-2, IL-4, IL-7, IL-9, and IL-15 (1). IL-21 is produced by activated T follicular helper cells (Tfh), Th17 cells, and NKT cells (2-6). It exerts its biological effects through a heterodimeric receptor complex of γ_c and the IL-21-specific IL-21 R (2, 7). Tfh-derived IL-21 plays an important role in the development of humoral immunity through its autocrine effects on the Tfh cell and paracrine effects on immunoglobulin affinity maturation, plasma cell differentiation, and B cell memory responses (4, 8, 9). It is also required for the migration of dendritic cells to draining lymph nodes (10). IL-21 regulates several aspects of T cell function. It co-stimulates the activation, proliferation, and survival of CD8+ T cells and NKT cells and promotes Th17 cell polarization (3, 5, 6, 11, 12). It blocks the generation of regulatory T cells and their suppressive effects on CD4+ T cells (13, 14). IL-21 R engagement enhances the cytolytic activity and IFN- γ production of activated NK cells but limits the expansion of resting NK cells (15). In addition, IL-21 suppresses cutaneous hypersensitivity reactions by limiting allergen-specific IgE production and mast cell degranulation (16). Dysregulation of the IL-21/IL-21 R system contributes to the development of multiple immunological disorders (1, 17). The 133 amino acid (aa) mature human IL-21 shares 63% and 61% aa sequence identity with mouse and rat IL-21, respectively. Alternative splicing generates an additional isoform with a substitution of the C-terminal 16 amino acids (18).

References:

1. Tangye, S.G. (2015) *Curr. Opin. Immunol.* **34**:107.
2. Parrish-Novak, *et al.* (2000) *Nature* **408**:57.
3. Coquet, J.M. *et al.* (2007) *J. Immunol.* **178**:2827.
4. Vogelzang, A. *et al.* (2008) *Immunity* **29**:127.
5. Korn, T. *et al.* (2007) *Nature* **448**:484.
6. Nurieva, R. *et al.* (2007) *Nature* **448**:480.
7. Asao, H. *et al.* (2001) *J. Immunol.* **167**:1.
8. Zotos, D. *et al.* (2010) *J. Exp. Med.* **207**:365.
9. Rankin, A.L. *et al.* (2011) *J. Immunol.* **186**:667.
10. Jin, H. *et al.* (2009) *J. Clin. Invest.* **119**:47.
11. Frohlich, A. *et al.* (2009) *Science* **324**:1576.
12. Yi, J.S., *et al.* (2009) *Science* **324**:1572.
13. Peluso, I. *et al.* (2007) *J. Immunol.* **178**:732.
14. Bucher, C. *et al.* (2009) *Blood* **114**:5375.
15. Kasaian, M.T. *et al.* (2002) *Immunity* **16**:559.
16. Tamagawa-Mineoka, R. *et al.* (2011) *J. Invest. Dermatol.* **131**:1513.
17. Ma, J. *et al.* (2011) *Cytokine* **56**:133.
18. Rahman, M. *et al.* (2007) *FEBS Lett.* **581**:4001.

MANUFACTURING SPECIFICATIONS

Animal-Free Manufacturing Conditions

Our dedicated controlled-access animal-free laboratories ensure that at no point in production are the products exposed to potential contamination by animal components or byproducts. Every stage of manufacturing is conducted in compliance with R&D Systems' stringent Standard Operating Procedures (SOPs). Production and purification procedures use equipment and media that are confirmed animal-free.

Production

- All molecular biology procedures use animal-free media and dedicated labware.
- Dedicated fermentors are utilized in committed animal-free areas.

Purification

- Protein purification columns are animal-free.
- Bulk proteins are filtered using animal-free filters.
- Purified proteins are stored in animal-free containers in a dedicated cold storage room.

Quality Assurance

- Low Endotoxin Level.
- No impairment of biological activity.
- High quality product obtained under stringent conditions.
- For *ex vivo* research or bioproduction, [additional documentation](#) can be provided.

[Please read our complete Animal-Free Statement.](#)