

DESCRIPTION

Source	Human embryonic kidney cell, HEK293-derived sars-cov-2 Spike protein			
	<p>SARS-CoV-2 C.37 Spike (Val16-Lys1211) (Gly75Val, Thr76Ile, RSYLTPG246-252 del, Asp253Asn, Leu452Gln, Phe490Ser, Asp614Gly, Thr859Asn) (Arg682Ser, Arg685Ser, Lys986Pro, Val987Pro) Accession # YP_009724390.1</p>	GCN4-IZ	Avi-tag	6-His tag
	N-terminus			C-terminus
N-terminal Sequence Analysis	Val16			
Structure / Form	Biotinylated via Avi-tag.			
Predicted Molecular Mass	137 kDa			

SPECIFICATIONS


SDS-PAGE	145 - 170 kDa, under reducing conditions.
Activity	Measured by its binding ability in a functional ELISA with Recombinant Human ACE-2 Fc Chimera (Catalog # 10544-ZN).
Endotoxin Level	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 500 µg/mL in PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after reconstitution.

DATA


Binding Activity



Biotinylated Recombinant SARS-CoV-2 C.37 Spike (GCN4-IZ) Avi-tag His-tag (ng/mL)

Biotinylated Recombinant SARS-CoV-2 C.37 Spike (GCN4-IZ) Avi-tag His-tag Protein Binding Activity.
Biotinylated Recombinant SARS-CoV-2 C.37 Spike (GCN4-IZ) Avi-tag His-tag (Catalog # AVI10887) binds Recombinant Human ACE-2 Fc Chimera (Catalog # 10544-ZN) in a functional ELISA.

SDS-PAGE



Biotinylated Recombinant SARS-CoV-2 C.37 Spike (GCN4-IZ) Avi-tag His-tag Protein SDS-PAGE. 2 µg/lane of Biotinylated Recombinant SARS-CoV-2 C.37 Spike (GCN4-IZ) Avi-tag His-tag Protein (Catalog # AVI10887) was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining, showing bands at 145-170 kDa.

BACKGROUND

SARS-CoV-2, which causes the global pandemic coronavirus disease 2019 (Covid-19), belongs to a family of viruses known as coronaviruses that also include MERS-CoV and SARS-CoV-1. Coronaviruses are commonly comprised of four structural proteins: Spike protein (S), Envelope protein (E), Membrane protein (M) and Nucleocapsid protein (N) (1). The SARS-CoV-2 S protein is a glycoprotein that mediates membrane fusion and viral entry. The S protein is homotrimeric, with each ~180-kDa monomer consisting of two subunits, S1 and S2 (2). In SARS-CoV-2, as with most coronaviruses, proteolytic cleavage of the S protein into S1 and S2 subunits is required for activation. The S1 subunit is focused on attachment of the protein to the host receptor while the S2 subunit is involved with cell fusion (3-5). The S protein of SARS-CoV-2 shares 75% and 29% aa sequence identity with S protein of SARS-CoV-1 and MERS, respectively. The S Protein of the SARS-CoV-2 virus, like the SARS-CoV-1 counterpart, binds a metalloproteinase, Angiotensin-Converting Enzyme 2 (ACE-2), but with much higher affinity and faster binding kinetics through the receptor binding domain (RBD) located in the C-terminal region of S1 subunit (6). It has been demonstrated that the S Protein can invade host cells through the CD147/EMMPRIN receptor and mediate membrane fusion (7, 8). Polyclonal antibodies to the RBD of the SARS-CoV-2 protein have been shown to inhibit interaction with the ACE-2 receptor, confirming RBD as an attractive target for vaccinations or antiviral therapy (9). There is also promising work showing that the RBD may be used to detect presence of neutralizing antibodies present in a patient's bloodstream, consistent with developed immunity after exposure to the SARS-CoV-2 (10). Several emerging SARS-CoV-2 genomes have been identified with mutations compared to the Wuhan-Hu-1 SARS-CoV-2 reference sequence. First identified in South America in late 2020, the C.37, or Lambda, variant is considered a Variant of Interest (VOI) as it contains several mutations in the RBD domain that potentially affect viral fitness and transmissibility: L452Q and F490S (11). The F490S mutation, along with several mutations at position L452, has been associated with resistance to neutralization by multiple monoclonal antibodies (12). Our Avi-tag Biotinylated SARS-CoV-2 C.37 Spike (GCN4-IZ) His-tag protein features biotinylation at a single site contained within the Avi-tag, a unique 15 amino acid peptide. Protein orientation will be uniform when bound to streptavidin-coated surface due to the precise control of biotinylation and the rest of the protein is unchanged so there is no interference in the protein's bioactivity.

References:

1. Wu, F. *et al.* (2020) *Nature* **579**:265.
2. Tortorici, M.A. and D. Veerler (2019). *Adv. Virus Res.* **105**:93.
3. Bosch, B.J. *et al.* (2003). *J. Virol.* **77**:8801.
4. Belouzard, S. *et al.* (2009) *Proc. Natl. Acad. Sci.* **106**:5871.
5. Millet, J.K. and G. R. Whittaker (2015) *Virus Res.* **202**:120.
6. Ortega, J.T. *et al.* (2020) *EXCLI J.* **19**:410.
7. Wang, K. *et al.* (2020) *bioRxiv* <https://www.biorxiv.org/content/10.1101/2020.03.14.988345v1>.
8. Isabel, *et al.* (2020) *Sci Rep* **10**, 14031. <https://doi.org/10.1038/s41598-020-70827-z>.
9. Tai, W. *et al.* (2020) *Cell. Mol. Immunol.* <https://doi.org/10.1016/j.it.2020.03.007.1>.
10. Okba, N. M. A. *et al.* (2020). *Emerg. Infect. Dis.* <https://doi.org/10.3201/eid2607.200841>.
11. Romero, P.E. *et al.* (2021) *medRxiv* <https://doi.org/10.1101/2021.06.26.21259487>.
12. Liu, Z. *et al.* (2021) *Cell Host Microbe*. **29**:477.