

DESCRIPTION

Source	Human embryonic kidney cell, HEK293-derived human IL-1Rrp2/IL-1R6 protein			
	Human IL-1 R6 (Asp20-Tyr337) Accession # Q9HB29.2	DIEGRMD	Human IgG ₁ Fc (Pro100-Lys330)	Avi-tag
	N-terminus		C-terminus	
N-terminal Sequence Analysis	Protein identity confirmed by mass spectrometry.			
Structure / Form	Disulfide-linked homodimer Biotinylated via Avi-tag			
Predicted Molecular Mass	65 kDa (monomer)			

SPECIFICATIONS


SDS-PAGE	78-87 kDa, under reducing conditions.
Activity	Measured by its binding ability in a functional ELISA. When Recombinant Human IL-36β/IL-1F8 (Catalog # 6834-ILB/CF) is immobilized at 2 µg/mL (100 µL/well), Biotinylated Recombinant Human IL-1Rrp2/IL-1R6 Fc Chimera Avi-tag (Catalog # AV111116) binds with an ED ₅₀ of 1.50-15.0 µg/mL.
Endotoxin Level	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 500 µg/mL in PBS.
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after reconstitution.

DATA


Binding Activity



Biotinylated Recombinant Human IL-1Rrp2/IL-1R6 Fc Chimera Avi-tag (µg/mL)

Biotinylated Recombinant Human IL-1Rrp2/IL-1R6 Fc Chimera Avi-tag Protein Binding Activity. When Recombinant Human IL-36β/IL-1F8 (Catalog # 6834-ILB/CF) is immobilized at 2 µg/mL (100 µL/well), Biotinylated Recombinant Human IL-1Rrp2/IL-1R6 Fc Chimera Avi-tag Protein (Catalog # AV111116) binds with an ED₅₀ of 1.50-15.0 µg/mL.

SDS-PAGE



Biotinylated Recombinant Human IL-1Rrp2/IL-1R6 Fc Chimera Avi-tag Protein SDS-PAGE. 2 µg/lane of Biotinylated Recombinant Human IL-1Rrp2/IL-1R6 Fc Chimera Avi-tag Protein (Catalog # AV111116) was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining, showing bands at 78-87 kDa and 156-174 kDa, respectively.

BACKGROUND

The Interleukin 1 receptor family (IL-1 R) comprises at least eleven members including IL-1 RI (IL-1 R1), IL-1 RII (IL-1 R2), IL-1 RAcP (IL-1 R3), ST2 (T1/IL-1 R4), IL-18 Ra (IL-1 Rrp/IL-1 R5), IL-1Rrp2 (IL-1 RL2/IL-1R6), IL-18 Rb (AcPL/IL-1 R7), IL-1RAPL-1 (TIGIRR-2/IL-1 R8), and TIGIRR-1 (IL-1 R9) (1). All family members possess three immunoglobulin (Ig)-like domains in their extracellular region. Most members also have an intracellular TIR (Toll-like receptor/IL-1 receptor signaling) domain that is also conserved in the Toll-like receptor family. Related proteins, SIGIRR (single Ig domain-containing IL-1 R-related molecule) and IL-18BP, differ from the other members by having only one Ig domain (1). Human IL-1Rrp2 cDNA encodes a 561 amino acid (aa) residue precursor protein with a putative 19 aa signal peptide and a 318 aa extracellular domain. It shares 67% and 65% amino acid sequence identity with rat and mouse IL 1Rrp2, respectively. IL-1Rrp2 is expressed in lung epithelium, brain vasculature, kidney, testis, monocytes, skin-derived keratinocytes, fibroblasts and, to a lesser extent, endothelial cells (2, 3). IL-1Rrp2 has been shown to mediate the activation of the transcription factor NFκB by the IL-1 family ligands IL-1 F6, F8 or F9 (also known as IL-1ε), with IL-1RAcP as a cofactor (3, 4). Response to IL-1F9 is specifically antagonized by IL-1 F5 (also known as IL-1δ), an IL-1 family ligand that is most closely related to IL-1ra (3). IL-1Rrp2, IL-1 F5, and IL-1F9 are all up-regulated in lesional psoriasis skin, suggesting that the IL-1Rrp2 mediated signaling pathway may take part in local inflammatory responses (3). Our Avi-tag Biotinylated IL-1Rrp2 features biotinylation at a single site contained within the Avi-tag, a unique 15 amino acid peptide. Protein orientation will be uniform when bound to streptavidin-coated surface due to the precise control of biotinylation and the rest of the protein is unchanged so there is no interference in the protein's bioactivity

References:

1. Boraschi, D. and A. Tagliabue (2006) Vitam. Horm. **74**:229.
2. Lovenberg, T. W. *et al.* (1996) J. Neuroimmunol. **70**:113.
3. Debets, R. *et al.* (2001) J. Immunol. **167**:1440.
4. Towne, J. E. *et al.* (2004) J. Biol. Chem. **279**:13677.