biotechne

Biotinylated Recombinant Human SIRPγ/CD172g Fc Chimera Avi-tag

RDSYSTEMS

Catalog Number: AVI4486

DESCRIPTION				
Source	Chinese Hamster Ovary cell line, CHO-derived human SIRP gamma/CD172g protein			
	Human SIRPG (Glu29-Pro360) Accession # Q9P1W8.3	GGIEGRMD	Human IgG ₁ (Pro100-Lys330)	Avi-tag
	N-terminus C-terminus			
N-terminal Sequence Analysis	Glu29			
Structure / Form	Disulfide-linked homodimer Biotinylated via Avi-tag			
Predicted Molecular Mass	65 kDa			

SPECIFICATIONS			
SDS-PAGE	72-80 kDa, under reducing conditions.		
Activity	Measured by its binding ability in a functional ELISA. Biotinylated Recombinant Human SIRPγ/CD172g Fc Chimera Avi-tag (Catalog # AVI4486) binds Recombinant Human CD47 Fc Chimera (Catalog # 4670-CD) with an ED ₅₀ of 65.0-650 ng/mL.		
Endotoxin Level	<0.10 EU per 1 μ g of the protein by the LAL method.		
Purity	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.		
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.		

PREPARATION AND STORAGE Reconstitute at 500 ug/ml

DATA

Reconstitution	Reconstitute at 500 µg/mL in PBS.			
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. Use a manual defrost freezer and avoid repeated freeze-thaw cycles.			
Stability & Storage				
	 12 months from date of receipt, -20 to -70 °C as supplied. 			
	 1 month, 2 to 8 °C under sterile conditions after reconstitution. 			
	 3 months -20 to -70 °C under sterile conditions after reconstitution 			



Biotinylated Recombinant Human SIRPγ/CD172g Fc Chimera Avi-tag Protein Binding Activity. Biotinylated Recombinant Human SIRPγ/CD172g Fc Chimera Avitag Protein (Catalog # AVI4486) binds Recombinant Human CD47 Fc Chimera (Catalog # 4670-CD) with an ED₅₀ of 65.0-650 ng/mL.

SDS-PAGE



Biotinylated Recombinant Human SIRPy/CD172g Fc Chimera Avi-tag Protein SDS-PAGE. 2 µg/lane of Biotinylated Recombinant Human SIRPy/CD172g Fc Chimera Avitag Protein (Catalog # AVI4486) was resolved with SDS-PAGE under reducing (R) and nonreducing (NR) conditions and visualized by Coomassie® Blue staining, showing bands at 72-80 kDa and 140-160 kDa, respectively.

Rev. 8/15/2023 Page 1 of 2



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449

biotechne

RDsystems

Biotinylated Recombinant Human SIRPy/CD172g Fc Chimera Avi-tag

Catalog Number: AVI4486

BACKGROUND

Signal regulatory protein gamma (SIRP gamma, designated CD172g), also called SIRP beta 2, is a monomeric 45-47 kDa type I transmembrane protein belonging to the SIRP/SHPS (CD172) family of the Ig superfamily (1-5). SIRP members are "paired receptors" with homology in the extracellular domain but variability in the C-terminus and signaling function (1, 2). The 387 amino acid (aa) SIRP gamma sequence contains a 28 aa potential signal sequence, a 332 aa extracellular domain (ECD) with four potential N-glycosylation sites, a 23 aa transmembrane domain and a 4 aa cytoplasmic sequence. SIRP gamma contains one V-type Ig-like domain that contains a J-like sequence and two C1-type Ig-like domains within its ECD (1, 2). Isoforms that lack one (isoform 2, 276 aa) or two (isoform 3, 170 aa) membrane-proximal C-type Ig-like domains have been described (5). Within the ECD, human SIRP gamma isoform 1 shares 78% aa identity with human SIRP beta 1, and appears to have structurally similar orthologs only in rhesus monkey and chimpanzee (100% and 91% aa identity, respectively) (2). SIRP gamma is the only

SIRP known to be expressed on T cells, CD56^{bright} NK cells and activated NK cells; it is not expressed on myeloid cells (5, 6). It shows adhesion to CD47, but at lower affinity than SIRP alpha (6). Expression of SIRP gamma on T cells suggests a role as an accessory protein interacting with CD47-expressing antigen presenting cells (5, 6). Unlike SIRP alpha that has cytoplasmic ITIM domains, and SIRP beta 1 that interacts with DAP-12, SIRP gamma does not contain any obvious signaling motif (1, 2, 6). However, SIRP gamma -mediated adhesion appears to promote antigen-specific T cell proliferation and costimulate T cell activation (5). Engagement of CD47 by SIRP gamma was shown to induce apoptosis on T-cell and monocyte cell lines (6). Our Avi-tag Biotinylated Human SIRP gamma Fc Chimera protein features biotinylation at a single site contained within the Avi-tag, a unique 15 amino acid peptide. Protein orientation will be uniform when bound to streptavidin-coated surface due to the precise control of biotinylation and the rest of the protein is unchanged so there is no interference in the protein's bioactivity

References:

- 1. Barclay, A.N. & M.H. Brown (2006) Nat. Rev. Immunol. 6:457.
- 2. vanBeek, E.M. et al. (2005) J. Immunol. 175:7781.
- 3. van den Berg, T.K. et al. (2005) J. Immunol. 175:7788.
- 4. Ichigotani, Y. et al. (2000) J. Hum. Genet. 45:378.
- 5. Piccio, L. et al. (2005) Blood 105:2421.
- 6. Brooke, G. et al. (2004) J. Immunol. 173:2562.

Rev. 8/15/2023 Page 2 of 2



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449