

DESCRIPTION

Species Reactivity	Cotton Rat
Specificity	Detects cotton rat TNF- α in Western blots. In this format, greater than 50% cross-reactivity with recombinant mouse TNF- α is observed, 25% cross-reactivity with recombinant human TNF- α is observed and 5% cross-reactivity with recombinant porcine TNF- α is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant cotton rat TNF- α (R&D Systems, Catalog # 1011-CR) Leu1-Leu156 Accession # AAL18818
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 μ g/mL	Recombinant Cotton Rat TNF- α (Catalog # 1011-CR)

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Tumor necrosis factor alpha (TNF- α) also known as cachectin and TNFSF2, is the prototypic ligand of the TNF superfamily. It is a pleiotropic molecule that plays a central role in inflammation, apoptosis, and immune system development. TNF- α is produced by a wide variety of immune and epithelial cell types (1, 2). The 156 amino acid (aa) cotton rat TNF- α is homologous to a portion of the extracellular domain (ECD) of TNF- α from other species (3). It shares 64%-76% aa sequence identity with bovine, canine, equine, feline, human, mouse, porcine, rat, and rhesus TNF- α . The 26 kDa type 2 transmembrane protein is assembled intracellularly to form a noncovalently linked homotrimer (4). Ligation of this complex induces reverse signaling that promotes lymphocyte costimulation but diminishes monocyte responsiveness (5). Cleavage of membrane bound TNF- α by TACE/ADAM17 releases a 55 kDa soluble trimeric form of TNF- α (6, 7). TNF- α trimers bind the ubiquitous TNF RI and the hematopoietic cell-restricted TNF RII, both of which are also expressed as homotrimers (1, 8). TNF- α regulates lymphoid tissue development through control of apoptosis (2). It also promotes inflammatory responses by inducing the activation of vascular endothelial cells and macrophages (2). TNF- α is a key cytokine in the development of several inflammatory disorders (9). It contributes to the development of type 2 diabetes through its effects on insulin resistance and fatty acid metabolism (10, 11).

References:

1. Idriss, H.T. and J.H. Naismith (2000) *Microsc. Res. Tech.* **50**:184.
2. Hehlgans, T. and K. Pfeffer (2005) *Immunology* **115**:1.
3. Blanco, J.C. *et al.* (2004) *J. Interferon Cytokine Res.* **24**:21.
4. Tang, P. *et al.* (1996) *Biochemistry* **35**:8216.
5. Eissner G. *et al.* (2004) *Cytokine Growth Factor Rev.* **15**:353.
6. Black, R.A. *et al.* (1997) *Nature* **385**:729.
7. Moss, M.L. *et al.* (1997) *Nature* **385**:733.
8. Loetscher, H. *et al.* (1991) *J. Biol. Chem.* **266**:18324.
9. Clark, I.A. (2007) *Cytokine Growth Factor Rev.* **18**:335.
10. Romanatto, T. *et al.* (2007) *Peptides* **28**:1050.
11. Hector, J. *et al.* (2007) *Horm. Metab. Res.* **39**:250.