

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse IL-17 RB in ELISAs and Western blots. In sandwich immunoassays, less than 0.5% cross-reactivity with recombinant human IL-17 RB, recombinant mouse (rm) IL-17 and rmlIL-17 R is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse IL-17 RB Arg18-Gly286 Accession # Q9JIP3
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Mouse IL-17 RB Fc Chimera (Catalog # 1040-BR)
Mouse IL-17 RB Sandwich Immunoassay		Reagent
ELISA Capture	2-8 µg/mL	Mouse IL-17 RB Antibody (Catalog # MAB10401)
ELISA Detection	0.1-0.4 µg/mL	Mouse IL-17 RB Biotinylated Antibody (Catalog # BAF1040)
Standard		Recombinant Mouse IL-17 RB Fc Chimera (Catalog # 1040-BR)

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

The interleukin 17 (IL-17) family of cytokines, comprising six members (IL-17, IL-17B through IL-17F), are structurally related proteins with a conserved cysteine-knot structure. These proinflammatory cytokines can induce local cytokine production and are involved in the regulation of the immune response. The cognate receptors activated by some of these cytokines have been identified (1, 2). Interleukin-17B Receptor (IL-17RB), also known as IL-17 Rh1, IL-17E R and EVI27, represents the second receptor of the IL-17 family cytokines to be recognized (2-4). Mouse IL17B R cDNA encodes a 499 amino acid (aa) residue type I membrane protein with a putative 17 aa signal peptide, a 269 aa extracellular domain, a 21 aa transmembrane domain and a 192 aa cytoplasmic tail. As a result of alternative splicing, a secreted variant of IL-17B R also exists (4). Human and mouse IL-17B R share 76% aa sequence identity. The IL-17B R protein sequence is only 19.2% and 18.2% identical to the human and mouse IL-17 R sequence, respectively. However, the receptors share many conserved cysteine residues within their extracellular domains as well as additional conserved elements within their cytoplasmic domains. At least three additional type I transmembrane receptors with homology to IL-17 R, including IL-17 RL (IL-17 RC), IL-17 RD, and IL-17 RE, have been reported (2, 6). By Northern blot analysis, mouse IL-17B R is highly expressed in liver and testes and is expressed at lower levels in kidney and lung. It is also expressed in some hematopoietic cell lines, including selected T cell, B cell, and myeloid cell lines (2-4). The expression of IL-17B R is significantly up-regulated under inflammatory conditions (7). IL-17 RB binds strongly to IL-17E and weakly to IL-17B. It does not bind IL-17 or IL-17F. Activation of IL-17B R by its ligands results in the activation of nuclear factor kappa-B (2-4).

References:

1. Aggarwal, S. and A.L. Gurney (2002) J. Leukoc. Biol. **71**:1.
2. Moseley, T.A. *et al.* (2003) Cytokine & Growth Factor Rev. **14**:155.
3. Shi, Y. *et al.* (2000) J. Biol. Chem. **275**:19167.
4. Lee, J. *et al.* (2001) J. Biol. Chem. **276**:1660.
5. Tian, E. *et al.* (2000) Oncogene **19**:2098.
6. Haudenschild, D. *et al.* (2002) J. Biol. Chem. **277**:4309.
7. Hurst, S.D. *et al.* (2002) J. Immunol. **169**:443.