

ORDERING INFORMATION

Catalog Number: BAF1132

Lot Number: UTW01

Size: 50 μg

Formulation: 0.2 um filtered solution in PBS

with BSA

Storage: -20° C

Reconstitution: sterile 0.1% BSA in TBS

Specificity: mouse Netrin-4

Immunogen: NS0-derived rmNetrin-4

Ig Type: goat IgG

Applications: Western blot

Immunohistochemistry

Biotinylated Anti-mouse Netrin-4 Antibody

Preparation

Produced in goats immunized with purified, NS0-derived, recombinant mouse Netrin-4 (rmNetrin-4). Mouse Netrin-4 specific IgG was purified by mouse Netrin-4 affinity chromatography and then biotinylated.

Formulation

Lyophilized from a 0.2 μm filtered solution in phosphate-buffered saline (PBS) containing 50 μg of bovine serum albumin (BSA) per 1 μg of antibody.

Reconstitution

Reconstitute with sterile Tris-buffered saline pH 7.3 (20 mM Trizma base, 150 mM NaCl) containing 0.1% BSA. If 1 mL of buffer is used, the antibody concentration will be 50 μ g/mL.

Storage

Lyophilized samples are stable for twelve months from date of receipt when stored at -20° C to -70° C. Upon reconstitution, the antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Reconstituted antibody can also be aliquotted and stored frozen at -20° C to -70° C in a manual defrost freezer for six months without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Specificity

This antibody has been selected for use as a detection antibody in the applications listed below.

Applications

Western Blot - This antibody can be used at 0.1 - 0.2 μg/mL with the appropriate secondary reagents to detect mouse Netrin-4. The detection limit for rmNetrin-4 is approximately 5 ng/lane under non-reducing and reducing conditions. In this format, this antibody shows approximately 25% cross-reactivity with rhNetrin-4 and less than 1% cross-reactivity with rmNetrin-1, rchNetrin-2 and rmNetrin-G1a.

Immunohistochemistry - This antibody will detect Netrin-4 in cells and tissues. The working dilution is 5 - 15 μ g/mL. For chromogenic detection of labeling, use R&D Systems' Cell and Tissue Staining Kits (CTS Series).

Optimal dilutions should be determined by each laboratory for each application.

This product or the use of this product is covered by U.S. Patents owned by The Regents of the University of California. This product is for research use only and is not to be used for commercial purposes. Use of this product to produce products for sale or for diagnostic, therapeutic or drug discovery purposes is prohibited. In order to obtain a license to use this product for such purposes, contact The Regents of the University of California.

U.S. Patent # 5,565,331, 6,096,866, 6,017,714, 6,309,638, 6,670,451, and other U.S. and international patents pending.