



Biotinylated Anti-human Netrin-4 Antibody

ORDERING INFORMATION

Catalog Number: BAF1254

Lot Number: JCV01

Size: 50 µg

Formulation: 0.2 µm filtered solution in PBS
with BSA

Storage: -20° C

Reconstitution: sterile 0.1% BSA in TBS

Specificity: human Netrin-4

Immunogen: NS0-derived rhNetrin-4

Ig Type: goat IgG

Applications: Western blot
Immunohistochemistry

Preparation

Produced in goats immunized with purified, NS0-derived, recombinant human Netrin-4 (rhNetrin-4). Human Netrin-4 specific IgG was purified by human Netrin-4 affinity chromatography and then biotinylated.

Formulation

Lyophilized from a 0.2 µm filtered solution in phosphate-buffered saline (PBS) containing 50 µg of bovine serum albumin (BSA) per 1 µg of antibody.

Reconstitution

Reconstitute with sterile Tris-buffered saline pH 7.3 (20 mM Trizma base, 150 mM NaCl) containing 0.1% BSA. If 1 mL of buffer is used, the antibody concentration will be 50 µg/mL.

Storage

Lyophilized samples are stable for twelve months from date of receipt when stored at -20° C to -70° C. Upon reconstitution, the antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Reconstituted antibody can also be aliquotted and stored frozen at -20° C to -70° C **in a manual defrost freezer** for six months without detectable loss of activity. **Avoid repeated freeze-thaw cycles.**

Specificity

This antibody has been selected for use as a detection antibody in human Netrin-4 western blots.

Applications

Western Blot - This antibody can be used at 0.1 - 0.2 µg/mL with the appropriate secondary reagents to detect human Netrin-4. The detection limit for rhNetrin-4 is approximately 5 ng/lane and 25 ng/lane under non-reducing and reducing conditions, respectively. In this format, this antibody shows approximately 50% cross-reactivity with rmNetrin-4 and less than 2% cross-reactivity with rchNetrin-1, rchNetrin-2, rmNetrin-1 and rmNetrin-G1a.

Immunohistochemistry - This antibody will detect Netrin-4 in cells and tissues. The working dilution is 15 µg/mL. For chromogenic detection of labeling, use R&D Systems' Cell and Tissue Staining Kits (CTS Series).

Optimal dilutions should be determined by each laboratory for each application.

This product or the use of this product is covered by U.S. Patents owned by The Regents of the University of California. This product is for research use only and is not to be used for commercial purposes. Use of this product to produce products for sale or for diagnostic, therapeutic or drug discovery purposes is prohibited. In order to obtain a license to use this product for such purposes, contact The Regents of the University of California.

U.S. Patent # 5,824,775, 6,218,526, and other U.S. and international patents pending.

FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

R&D Systems, Inc.
1-800-343-7475