

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse SIGNR4/CD209e in Western blots. In Western blots, approximately 20% cross-reactivity with recombinant mouse SIGNR1 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse SIGNR4/CD209e Ser41-Trp208 Accession # Q91ZW7
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Mouse SIGNR4/CD209e

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Mouse SIGNR4 is a type II transmembrane C-type lectin that is homologous to human DC-SIGNR/DC-SIGN2. Mouse SIGNR4 is expressed at high levels in testis and at very low levels in the spleen (1).

References:

1. Park, C-G. *et al.* (2001) *Int. Immunol.* **13**:1283.