

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse B7-H2 in Western blots. In Western blots, less than 1% cross-reactivity with recombinant human (rh) B7-H2, recombinant mouse (rm) B7-1, rmB7-2, and rhB7-H1 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse B7-H2 Glu47-Lys279 Accession # Q9JHJ8
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Mouse B7-H2 Fc Chimera (Catalog # 158-B7)

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Mouse B7-H2, also called B7RP-1, B7h, LICOS, and GL50, is a member of the growing B7 family of immune costimulatory proteins. Other family members include B7-1, B7-2, B7-H1 (PD-L1), PD-L2, and B7-H3. B7 proteins are members of the immunoglobulin (Ig) superfamily. The extracellular domains contain 2 Ig-like domains and all members have short cytoplasmic domains. Among the family members, they share about 20-25% amino acid identity. Mouse and human B7-H2 share approximately 49% amino acid identity. B7-H2 has been identified as the ligand for ICOS, a member of the CD28 family of costimulatory receptors. Mouse B7-H2 is a 322 amino acid (aa) protein with a putative 46 aa signal peptide, a 233 aa extracellular domain, a 19 aa transmembrane region, and a 24 aa cytoplasmic domain. Mouse B7-H2 is expressed constitutively on resting B cells and at low levels on monocytes. The B7-H2/ICOS interaction appears to play roles in T cell dependent B cell activation and T_H differentiation.

References:

1. Coyle, A.J. and J.C. Gutierrez-Ramos (2001) *Nat. Immunol.* **2**:203.
2. Ling, V. *et al.* (2000) *J. Immunol.* **164**:1653.
3. Wang, S. *et al.* (2000) *Blood* **96**:2808.
4. Brodie, D. *et al.* (2000) *Curr. Biol.* **10**:333.
5. Mages, H.W. *et al.* (2000) *Eur. J. Immunol.* **30**:1040.
6. Swallow, M.M. *et al.* (1999) *Immunity* **11**:423.
7. Yoshinaga, S.K. *et al.* (1999) *Nature* **402**:827.