

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Otx2 in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Otx2 Met1-Leu289 Accession # P32243
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human Otx2
Immunocytochemistry	5-15 µg/mL	See Below

DATA

Immunocytochemistry



Otx2 in BG01V Human Embryonic Stem Cells. Otx2 was detected in immersion fixed BG01V human embryonic stem cells differentiated to ectoderm using Goat Anti-Human Otx2 Biotinylated Antigen Affinity-purified Polyclonal Antibody (Catalog # BAF1979) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Streptavidin (red; Catalog # NL999) and counterstained with DAPI (blue). Specific staining was localized to nuclei. Cells were co-stained using anti-Tubulin (Catalog # NB600-506, Novus Biologicals) and NorthernLights™ 493-conjugated Anti-Rat IgG Secondary Antibody (green; Catalog # NL015). View our protocol for [Fluorescent ICC Staining of Cells on Coverslips](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Otx2 is a member of the bicoid subfamily of homeodomain-containing transcription factors. It may play a role in brain and sensory organ development (1). Isoform a differs from isoform b by having an eight amino acid residue insertion between P32 and A33 of Otx2b.

References:

1. Nagao, T. *et al.* (1998) Proc. Natl. Acad. Sci. USA **95**:3737.