

DESCRIPTION



Species Reactivity	Human
Specificity	Detects human IL-1 β /IL-1F2 in ELISAs and Western blots. In sandwich immunoassays, less than 4% cross-reactivity with recombinant rat IL-1 β is observed and less than 0.1% cross-reactivity with recombinant mouse IL-1 β and recombinant porcine IL-1 β is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human IL-1 β /IL-1F2 Ala117-Ser269 Accession # NP_000567
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 μ g/mL	Recombinant Human IL-1 β /IL-1F2 (Catalog # 201-LB)
Immunocytochemistry	5-15 μ g/mL	See Below
Human IL-1β/IL-1F2 Sandwich Immunoassay		Reagent
ELISA Capture	2-8 μ g/mL	Human IL-1 β /IL-1F2 Antibody (Catalog # MAB601)
ELISA Capture	2-8 μ g/mL	Human IL-1 β /IL-1F2 Antibody (Catalog # MAB601R)
ELISA Detection	0.1-0.4 μ g/mL	Human IL-1 β /IL-1F2 Biotinylated Antibody (Catalog # BAF201)
Standard		Recombinant Human IL-1 β /IL-1F2 (Catalog # 201-LB)

DATA

Immunocytochemistry	IL-1 β /IL-1F2 in Human PBMCs. IL-1 β /IL-1F2 was detected in immersion fixed human peripheral blood mononuclear cells (PBMCs) using Human IL-1 β /IL-1F2 Biotinylated Antigen Affinity-purified Polyclonal Antibody (Catalog # BAF201) at 10 μ g/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Streptavidin (red; Catalog # NL999) and counterstained with DAPI (blue). View our protocol for Fluorescent ICC Staining of Non-adherent Cells .
	

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

IL-1 is a name that designates two pleiotropic cytokines, IL-1 α (IL-1F1) and IL-1 β (IL-1F2), which are the products of distinct genes. IL-1 α and IL-1 β are structurally related polypeptides that share approximately 21% amino acid (aa) identity in human. Both proteins are produced by a wide variety of cells in response to inflammatory agents, infections, or microbial endotoxins. While IL-1 α and IL-1 β are regulated independently, they bind to the same receptor and exert identical biological effects. IL-1 RI binds directly to IL-1 α or IL-1 β and then associates with IL-1 R accessory protein (IL-1 R3/IL-1 R AcP) to form a high-affinity receptor complex that is competent for signal transduction. IL-1 RII has high affinity for IL-1 β but functions as a decoy receptor and negative regulator of IL-1 β activity. IL-1ra functions as a competitive antagonist by preventing IL-1 α and IL-1 β from interacting with IL-1 RI. The human IL-1 β cDNA encodes a 269 aa precursor. A 116 aa propeptide is cleaved intracellularly by the cysteine protease IL-1 β -converting enzyme (Caspase-1/ICE) to generate the active cytokine. The 17 kDa mature human IL-1 β shares 96% aa sequence identity with rhesus and 67-78% with canine, cotton rat, equine, feline, mouse, porcine, and rat IL-1 β .