

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-7 in ELISAs and Western blots. In sandwich immunoassays, less than 0.05% cross-reactivity with recombinant mouse (rm) IL-7, rmlIL-7 Ra, and rhIL-7 Ra is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human IL-7 (R&D Systems, Catalog # 207-IL) Asp26-His177 Accession # P13232
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human IL-7 (Catalog # 207-IL)
Human IL-7 Sandwich Immunoassay		Reagent
ELISA Capture	2-8 µg/mL	Human IL-7 Antibody (Catalog # MAB207)
ELISA Detection	0.1-0.4 µg/mL	Human IL-7 Biotinylated Antibody (Catalog # BAF207)
Standard		Recombinant Human IL-7 (Catalog # 207-IL)

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

IL-7, previously known as pre-B-cell growth factor and lymphopoietin-1, was originally purified on the basis of its ability to promote the proliferation of precursor B-cells. It has now been shown that IL-7 can also stimulate the proliferation of thymocytes, T cell progenitors and mature CD4⁺ and CD8⁺ T cells. IL-7 can induce the formation of lymphokine-activated killer (LAK) cells as well as the development of cytotoxic T lymphocytes (CTL). IL-7 was also shown to induce the V(D)J rearrangement of the T cell receptor β gene in mouse fetal thymocytes. Among myeloid lineage cells, IL-7 can up-regulate the production of pro-inflammatory cytokines and stimulate the tumoricidal activity of monocytes/macrophages. IL-7 is expressed by adherent stromal cells from various tissues.

Human IL-7 cDNA encodes a precursor protein of 177 amino residues containing a 25 amino acid residue signal peptide. Mouse IL-7 has approximately 65% amino acid sequence identity with human IL-7 and both proteins exhibit cross-species activity. IL-7 bioactivities are mediated by the binding of IL-7 to functional high-affinity receptor complexes. The ligand binding subunit (IL-7 R) of the IL-7 receptor complex has been cloned from human and mouse sources. In addition to the membrane-anchored form of the IL-7 receptor, a human cDNA clone that encodes a soluble form of the IL-7 R has also been isolated. The γ chain of the IL-2 receptor complex has been shown to be an essential component for IL-7 signal transduction. Both IL-7 R and IL-2 R_γ are members of the hematopoietin receptor superfamily. Cells known to express IL-7 receptors include pre-B cells, T cells and bone marrow cells.