

Human SIRP beta 1/gamma Biotinylated Antibody

Antigen Affinity-purified Polyclonal Goat IgG Catalog Number: BAF2096

DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human SIRP beta 1 and human SIRP gamma in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human SIRP beta 1 Gly26-Ala369 (Arg53His and Ala363Pro) Accession # 000241
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS				
Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.				
	Recommended	Sample		
	Concentration			
Western Blot	0.1 μg/mL	Recombinant Human SIRP beta 1 and Recombinant Human SIRP gamma		

PREPARATION AND STORAGE			
Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.		
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.		
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.		

BACKGROUND

SIRP\$\beta\$1 is a type I transmembrane protein belonging to the SIRP family within the Ig superfamily. Members of this family are characterized by an extracellular region containing a V-set Ig domain containing a J-like sequence and two C1-set Ig domains. Unlike SIRP\$\alpha\$ that has cytoplasmic ITIM domains, SIRP\$\beta\$1 possesses positively charged residues that allow association with ITAM motif containing adaptor molecules. SIRP\$\beta\$1 is expressed on cells of monocyte, macrophage or dendritic lineages.

