

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse NKp46 in Western blots. In this format, approximately 10% cross-reactivity with recombinant human (rh) NKp46 is observed and less than 1% cross-reactivity with rhNKp30 and rhNKp80 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse NKp46 Glu22-Asn255 Accession # Q8C567
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Mouse NKp46/NCR1 Fc Chimera (Catalog # 2225-NK)

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

NKp46, along with NKp30 and NKp44, are activating receptors that have been collectively termed the natural cytotoxicity receptors (NCR) (1). These receptors are expressed almost exclusively by NK cells and play a major role in triggering some of the key lytic activities of NK cells. In human systems, the CD56^{dim}CD16⁺ subpopulation that makes up the majority of NK cells in the peripheral blood and spleen expresses NKp46 in both resting and activated states (2). The main NK cell population of the lymph node (CD56^{bright}CD16⁻) expresses low levels of NKp46 in resting cells, but expression is upregulated by IL-2. Mouse NKp46, also known as MAR-1 (3), is a type I transmembrane protein with two extracellular Ig-like domains. It has a positive charge in its transmembrane domain that permits association with the ITAM-bearing signal adapter proteins, CD3 ζ and Fc ϵ R1 γ (4). Studies with neutralizing antibodies indicate that the three NCR are primarily responsible for triggering the NK-mediated lysis of many human tumor cell lines. Blocking any of the NCRs individually resulted in partial inhibition of tumor cell lysis, but nearly complete inhibition of lysis was observed if all three receptors were blocked simultaneously (5). NKp46 has also been implicated in recognition of virus-infected cells through its capacity to bind to viral hemagglutinins (6 - 8).

References:

1. Moretta, L. and A. Moretta (2004) EMBO J. **23**:255.
2. Ferlazzo, G. *et al.* (2004) J. Immunol. **172**:1455.
3. Biassoni, R. *et al.* (1999) Eur. J. Immunol. **29**:1014.
4. Westgaard, I. *et al.* (2004) J. Leukoc. Biol. PMID 15356098.
5. Pende, D. *et al.* (1999) J. Exp. Med. **190**:1505.
6. Arnon, T. *et al.* (2004) Blood **103**:664.
7. Arnon, T. *et al.* (2001) Eur. J. Immunol. **31**:2680.
8. Mandelboim, O. *et al.* (2001) Nature **409**:1055.