

## DESCRIPTION

<b>Species Reactivity</b>	Bovine
<b>Specificity</b>	Detects TNF- $\alpha$ in ELISAs and Western blots. In sandwich immunoassays, less than 0.3% cross-reactivity with recombinant human TNF- $\alpha$ , recombinant mouse TNF- $\alpha$ , recombinant rat TNF- $\alpha$ , recombinant feline TNF- $\alpha$ , recombinant equine TNF- $\alpha$ , recombinant porcine TNF- $\alpha$ , recombinant cotton rat TNF- $\alpha$ , and rhesus macaque TNF- $\alpha$ is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant bovine TNF- $\alpha$ Leu78-Leu234 Accession # Q06599
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.


## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	0.1 $\mu$ g/mL	Recombinant Bovine TNF- $\alpha$ (Catalog # 2279-BT)
<b>Immunocytochemistry</b>	5-15 $\mu$ g/mL	See Below
<b>Bovine TNF-<math>\alpha</math> Sandwich Immunoassay</b>		<b>Reagent</b>
<b>ELISA Capture</b>	0.2-0.8 $\mu$ g/mL	Bovine TNF- $\alpha$ Antibody (Catalog # AF2279)
<b>ELISA Detection</b>	0.1-0.4 $\mu$ g/mL	Bovine TNF- $\alpha$ Biotinylated Antibody (Catalog # BAF2279)
<b>Standard</b>		Recombinant Bovine TNF- $\alpha$ (Catalog # 2279-BT)

## DATA

### Immunocytochemistry



**TNF- $\alpha$  in Bovine PBMCs.** TNF- $\alpha$  was detected in immersion fixed bovine peripheral blood mononuclear cells (PBMCs) using Goat Anti-Bovine TNF- $\alpha$  Biotinylated Antigen Affinity-purified Polyclonal Antibody (Catalog # BAF2279) at 15  $\mu$ g/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Streptavidin (red; Catalog # NL999) and counterstained with DAPI (blue). Specific staining was localized to cytoplasm. View our protocol for [Fluorescent ICC Staining of Non-adherent Cells](#).

## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<p><b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b></p> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

Tumor necrosis factor alpha (TNF- $\alpha$ , also known as cachectin and TNFSF2, is the prototypic ligand of the TNF superfamily. It is a pleiotropic molecule that plays a central role in inflammation, apoptosis, and immune system development. TNF- $\alpha$  is produced by a wide variety of immune and epithelial cell types (1, 2). Bovine TNF- $\alpha$  consists of a 35 amino acid (aa) cytoplasmic domain, a 21 aa transmembrane segment, and a 178 aa extracellular domain (ECD) (3). Within the ECD, bovine TNF- $\alpha$  shares 64%-83% sequence identity with canine, cotton rat, equine, feline, human, mouse, porcine, rat, and rhesus TNF- $\alpha$ . The 26 kDa type 2 transmembrane protein is assembled intracellularly to form a noncovalently linked homotrimer (4). Ligation of this complex induces reverse signaling that promotes lymphocyte costimulation but diminishes monocyte responsiveness (5). Cleavage of membrane bound TNF- $\alpha$  by TACE/ADAM17 releases a 55 kDa soluble trimeric form of TNF- $\alpha$  (6, 7). TNF- $\alpha$  trimers bind the ubiquitous TNF RI and the hematopoietic cell-restricted TNF RII, both of which are also expressed as homotrimers (1, 8). TNF- $\alpha$  regulates lymphoid tissue development through control of apoptosis (2). It also promotes inflammatory responses by inducing the activation of vascular endothelial cells and macrophages (2). TNF- $\alpha$  is a key cytokine in several inflammatory disorders (9). It contributes to the development of type 2 diabetes through its effects on insulin resistance and fatty acid metabolism (10, 11).

**References:**

1. Idriss, H.T. and J.H. Naismith (2000) *Microsc. Res. Tech.* **50**:184.
2. Hehlgans, T. and K. Pfeffer (2005) *Immunology* **115**:1.
3. Cludts, I. *et al.* (1993) *Cytokine* **5**:336.
4. Tang, P. *et al.* (1996) *Biochemistry* **35**:8216.
5. Eissner G. *et al.* (2004) *Cytokine Growth Factor Rev.* **15**:353.
6. Black, R.A. *et al.* (1997) *Nature* **385**:729.
7. Moss, M.L. *et al.* (1997) *Nature* **385**:733.
8. Loetscher, H. *et al.* (1991) *J. Biol. Chem.* **266**:18324.
9. Clark, I.A. (2007) *Cytokine Growth Factor Rev.* **18**:335.
10. Romanatto, T. *et al.* (2007) *Peptides* **28**:1050.
11. Hector, J. *et al.* (2007) *Horm. Metab. Res.* **39**:250.