

DESCRIPTION

Species Reactivity	Human/Primate
Specificity	Detects human EGF in ELISAs and Western blots. In sandwich immunoassays, less than 0.01% cross-reactivity with rmEGF, rhEGF R and rhHB-EGF is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human EGF
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human EGF (Catalog # 236-EG) under non-reducing conditions only
Human/Primate EGF Sandwich Immunoassay		Reagent
ELISA Capture	2-8 µg/mL	Human/Primate EGF Antibody (Catalog # MAB636)
ELISA Detection Standard	0.1-0.4 µg/mL	Human/Primate EGF Biotinylated Antibody (Catalog # BAF236) Recombinant Human EGF (Catalog # 236-EG)

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Epidermal growth factor (EGF) is the founding member of the EGF family that also includes TGF-α, amphiregulin (AR), betacellulin (BTC), epiregulin (EPR), heparin-binding EGF-like growth factor (HB-EGF), epigen, and the neuregulins (NRG)-1 through -6 (1). Members of the EGF family share a structural motif, the EGF-like domain, which is characterized by three intramolecular disulfide bonds that are formed by six similarly spaced conserved cysteine residues (2). All EGF family members are synthesized as type I transmembrane precursor proteins that may contain several EGF domains in the extracellular region. The mature proteins are released from the cell surface by regulated proteolysis (1). The 1207 amino acid (aa) human EGF precursor contains nine EGF domains and nine LDLR class B repeats. The mature protein consists of 53 aa and is generated by proteolytic excision of the EGF domain proximal to the transmembrane region (3). Mature human EGF shares 70% aa sequence identity with mature mouse and rat EGF. EGF is present in various body fluids, including blood, milk, urine, saliva, seminal fluid, pancreatic juice, cerebrospinal fluid, and amniotic fluid (4). Four ErbB (HER) family receptor tyrosine kinases including EGFR/ErbB1, ErbB2, ErbB3 and ErbB4, mediate responses to EGF family members (5). These receptors undergo a complex pattern of ligand induced homo- or hetero-dimerization to transduce EGF family signals (6, 7). EGF binds ErbB1 and depending on the context, induces the formation of homodimers or heterodimers containing ErbB2. Dimerization results in autophosphorylation of the receptor at specific tyrosine residues to create docking sites for a variety of signaling molecules (5, 8). Biological activities ascribed to EGF include epithelial development, angiogenesis, inhibition of gastric acid secretion, fibroblast proliferation, and colony formation of epidermal cells in culture.

References:

1. Harris, R.C. *et al.* (2003) *Exp. Cell Res.* **284**:2.
2. Carpenter, G. and Cohen, S. (1990) *J. Biol. Chem.* **265**:7709.
3. Bell, G.I. *et al.* (1986) *Nucl. Acids Res.* **14**:8427.
4. Carpenter, G. and Zengdegi, J.G. (1986) *Exp. Cell Res.* **164**:1.
5. Jorissen, R.N. *et al.* (2003) *Exp. Cell Res.* **284**:31.
6. Gamett, D.C. *et al.* (1997) *J. Biol. Chem.* **272**:12052.
7. Qian, X. *et al.* (1994) *Proc. Natl. Acad. Sci.* **91**:1500.
8. Qian, X. *et al.* (1999) *J. Biol. Chem.* **274**:574.