

## DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects TREML1/TLT-1 in Western blots. In Western blots, less than 2% cross-reactivity with recombinant mouse TLT-1 is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human TREML1 Gln16-Ser160 (predicted) Accession # Q86YW5
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.1 µg/mL	Recombinant Human TREML1/TLT-1
<b>Flow Cytometry</b>	2.5 µg/10 <sup>6</sup> cells	Human platelets

## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<p><b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b></p> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

TREML1 is a type I transmembrane protein that differs from TREM in the cytoplasmic region by having two ITIMs (Immunoreceptor Tyrosine-based Inhibition Motif). A splice variant lacking ITIMs in its cytoplasmic domain also exists. TREML1 is found only in platelets and megakaryocytes. The extracellular region of human and mouse TREML1 share 75% amino acid sequence identity.