

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse MICL/CLEC12A in Western blots. In Western blots, less than 1% cross-reactivity with recombinant human MICL is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse MICL/CLEC12A Thr67-Arg267 Accession # NP_808354
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Mouse MICL/CLEC12A

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

MICL, also known as killer cell lectin-like receptor 1 (KLR1), is an inhibitory C-type lectin-like receptor that contains an immunoreceptor tyrosine-based inhibitory motif (ITIM). It is a 75 kDa type II membrane protein that is expressed on neutrophils, NK cells, monocytes and dendritic cells. The amino acid sequence of mouse MICL extracellular domain is 53%, 52% and 58% identical to that of human, bovine and canine MICL, respectively.