

## DESCRIPTION

<b>Species Reactivity</b>	Human/Mouse/Rat
<b>Specificity</b>	Detects Activin A in Western blots.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human Activin A
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	0.1 µg/mL	Recombinant Human/Mouse/Rat Activin A (Catalog # 338-AC)

## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<p><b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b></p> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Activin and Inhibin are members of the TGF-β superfamily of cytokines and are involved in a wide range of biological processes including tissue morphogenesis and repair, fibrosis, inflammation, neural development, hematopoiesis, reproductive system function, and carcinogenesis (1-7). Activin and Inhibin are produced as precursor proteins. Their amino terminal propeptides are proteolytically cleaved and facilitate formation of disulfide-linked dimers of the bioactive proteins (8, 9). Activins are nonglycosylated homodimers or heterodimers of various β subunits (βA, βB, βC, and βE in mammals), while Inhibins are heterodimers of a unique α subunit and one of the β subunits. Activin A is a widely expressed homodimer of two βA chains. The βA subunit can also heterodimerize with a βB or βC subunit to form Activin AB and Activin AC, respectively (10). The 14 kDa mature human βA chain shares 100% amino acid sequence identity with bovine, feline, mouse, porcine, and rat βA. Activin A exerts its biological activities by binding to the type 2 serine/threonine kinase Activin RIIA which then noncovalently associates with the type 1 serine/threonine kinase Activin RIB/ALK-4 (7, 11). Signaling through this receptor complex leads to Smad activation and regulation of activin-responsive gene transcription (7, 11). The bioactivity of Activin A is regulated by a variety of mechanisms (11). BAMBI, Betaglycan, and Cripto are cell-associated molecules that function as decoy receptors or limit the ability of Activin A to induce receptor complex assembly (12-14). The intracellular formation of Activin A can be prevented by the incorporation of the βA subunit into Activin AC or Inhibin A (3, 10). And the bioavailability of Activin A is restricted by its incorporation into inactive complexes with α2-Macroglobulin, Follistatin, and FLRG (15, 16).

### References:

1. Kumanov, P. *et al.* (2005) *Reprod. Biomed. Online* **10**:786.
2. Maeshima, A. *et al.* (2008) *Endocr. J.* **55**:1.
3. Rodgarkia-Dara, C. *et al.* (2006) *Mutat. Res.* **613**:123.
4. Werner, S. and C. Alzheimer (2006) *Cytokine Growth Factor Rev.* **17**:157.
5. Xu, P. and A.K. Hall (2006) *Dev. Biol.* **299**:303.
6. Shav-Tal, Y. and D. Zipori (2002) *Stem Cells* **20**:493.
7. Chen, Y.G. *et al.* (2006) *Exp. Biol. Med.* **231**:534.
8. Gray, A.M. and A.J. Mason (1990) *Science* **247**:1328.
9. Mason, A.J. *et al.* (1996) *Mol. Endocrinol.* **10**:1055.
10. Thompson, T.B. *et al.* (2004) *Mol. Cell. Endocrinol.* **225**:9.
11. Harrison, C.A. *et al.* (2005) *Trends Endocrinol. Metab.* **16**:73.
12. Onichtchouk, D. *et al.* (1999) *Nature* **401**:480.
13. Gray, P.C. *et al.* (2002) *Mol. Cell. Endocrinol.* **188**:254.
14. Kelber, J.A. *et al.* (2008) *J. Biol. Chem.* **283**:4490.
15. Phillips, D.J. *et al.* (1997) *J. Endocrinol.* **155**:65.
16. Schneyer, A. *et al.* (2003) *Endocrinology* **144**:1671.