

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse GLI-1 in Western blot. In Western blots, less than 1% cross-reactivity with recombinant mouse (rm) GLI-2 and rmGLI-3 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse GLI-1 Met1-Glu237 Accession # P47806
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human GLI-1 Recombinant Mouse GLI-1

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

GLI-1 is a member of the Kruppel family of zinc finger proteins. GLI-1 is activated by the sonic hedgehog pathway and influences transcription of a variety of target genes by binding to the consensus site 5'CGGGTGGTC3'. GLI-1 activation leads to cellular proliferation and anti-apoptotic activities based on target genes activated.