

## **Rat CD34 Biotinylated Antibody**

Antigen Affinity-purified Polyclonal Goat IgG Catalog Number: BAF4117

DESCRIPTION	
Species Reactivity	Rat
Specificity	Detects rat CD34 in Western blots. In Western blots, less than 1% cross-reactivity with recombinant porcine CD34 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant rat CD34 Glu37-Ile293 Accession # NP_001100672
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.
Please Note: Optimal diluti	ions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.  Recommended Sample Concentration
Western Blot	0.1 μg/mL Recombinant Rat CD34
PREPARATION AND S	STORAGE
Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.  • 12 months from date of receipt, -20 to -70 °C as supplied.

## BACKGROUND

Rat CD34 is a 80-100 kDa type I transmembrane glycoprotein that belongs to the CD34/Podocalyxin family of sialomucins. It is a widely used marker of activated hematopoietic stem/progenitor cells and is also expressed on adult vascular endothelial cells. Mature rat CD34 is predicted to be a 352 amino acid (aa) protein with a 257 aa extracellular domain (ECD) (aa 35-291) and 74 aa cytoplasmic tail (aa 313-386). Over the region used as immunogen, rat CD34 shares 83% and 57% aa identity with mouse and human CD34, respectively.

1 month, 2 to 8 °C under sterile conditions after reconstitution.
 6 months, -20 to -70 °C under sterile conditions after reconstitution.

biotechne