

ORDERING INFORMATION

Catalog Number: BAF4294

Lot Number: CAXI01

Size: 50 μg

Formulation: 0.2 um filtered solution in PBS

with BSA

Storage: -20° C

Reconstitution: sterile 0.1% BSA in TBS

Specificity: mouse IL-22 R

extracellular domain

Immunogen: NS0-derived rmIL-22 R

extracellular domain

Ig Type: sheep IgG

Application: Western blot

Biotinylated Anti-mouse IL-22 R Antibody

Preparation

Produced in sheep immunized with purified, NS0-derived, recombinant mouse IL-22 R extracellular domain (rmIL-22 R; R&D Systems, Catalog # 4294-MR). Mouse IL-22 R specific IgG was purified by mouse IL-22 R affinity chromatography and then biotinylated.

Formulation

Lyophilized from a 0.2 μ m filtered solution in phosphate-buffered saline (PBS) containing 50 μ g of bovine serum albumin (BSA) per 1 μ g of antibody.

Reconstitution

Reconstitute with sterile Tris-buffered saline pH 7.3 (20 mM Trizma base, 150 mM NaCl) containing 0.1% BSA. If 1 mL of buffer is used, the antibody concentration will be 50 μ g/mL.

Storage

Lyophilized samples are stable for twelve months from date of receipt when stored at -20° C to -70° C. Upon reconstitution, the antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Reconstituted antibody can also be aliquotted and stored frozen at -20° C to -70° C in a manual defrost freezer for six months without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Specificity

This antibody has been selected for use as a detection antibody in mouse IL-22 R Western blots.

Application

Western blot - This antibody can be used at 0.1 - 0.2 μ g/mL with the appropriate secondary reagents to detect mouse IL-22 R. The detection limit for rmIL-22 R is approximately 2 ng/lane under non-reducing and reducing conditions. In this format, this antibody shows approximately 30% cross-reactivity with rhIL-22 R.

Optimal dilutions should be determined by each laboratory for each application.