

#### DESCRIPTION

<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse CD40 in ELISAs and Western blots. In sandwich immunoassays, less than 0.05% cross-reactivity with rhCD40, rmCD27, rmCD30 and rmFas is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant mouse CD40 isoform 1 Val24-Arg193 Accession # P27512
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.1 µg/mL	Recombinant Mouse CD40/TNFRSF5 Fc Chimera (Catalog # <a href="#">1215-CD</a> )
<b>Mouse CD40/TNFRSF5 Sandwich Immunoassay</b>		<b>Reagent</b>
<b>ELISA Capture</b>	2-8 µg/mL	Mouse CD40/TNFRSF5 Antibody (Catalog # <a href="#">MAB4401</a> )
<b>ELISA Detection</b>	0.1-0.4 µg/mL	Mouse CD40/TNFRSF5 Biotinylated Antibody (Catalog # <a href="#">BAF440</a> )
<b>Standard</b>		Recombinant Mouse CD40/TNFRSF5 Fc Chimera (Catalog # <a href="#">1215-CD</a> )

#### PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

#### BACKGROUND

CD40 is a type I transmembrane glycoprotein belonging to the TNF receptor superfamily. The mature mCD40 consists of a 172 amino acid (aa) extracellular domain, a 22 aa transmembrane region and a 90 aa cytoplasmic domain (1). CD40 is expressed on B cells, follicular dendritic cells, dendritic cells, activated monocytes, macrophages, endothelial cells, vascular smooth muscle cells and several tumor cell lines (2). The extracellular domain has the cysteine-rich repeat regions, which are characteristic for many of the receptors of the TNF superfamily. Interaction of CD40 with its ligand, CD40L, leads to the aggregation of CD40 molecules, which in turn interact with cytoplasmic components to initiate signaling pathways. Early studies on the CD40-CD40L system revealed its role in humoral immunity. Interaction between CD40L on T cells and CD40 on B cells stimulated B cell proliferation and provided the signal for immunoglobulin isotype switching (3). Mutations in the CD40L gene, which resulted in a CD40L molecule unable to interact with CD40, are responsible for the hyper-IgM syndrome (4). Cross-linking of CD40 with antibodies or by binding to CD40L produces cell type-specific responses which include costimulation and induction of proliferation, induction of cytokine production, rescue from apoptosis, and upregulation of adhesion molecules (5). Some of the early events of intracellular signaling by the CD40-CD40L system include the association of the CD40 with TRAFs and the activation of various kinases (6 - 8).

#### References:

- Torres, R.M. and E.A. Clark (1992) J. Immunol. **148**:620.
- Schonbeck, U. *et al.* (1997) J. Biol. Chem. **272**:19569.
- Armitage, R.J. *et al.* (1993) J. Immunol. **150**:3671.
- Callard, R.E. *et al.* (1993) Immunol. Today **14**:559.
- Stout, R.D. and J. Suttles (1996) Immunol. Today **17**:487.
- Pullen, S.S. *et al.* (1999) Biochemistry **38**:10168.
- Faris, M. *et al.* (1994) J. Exp. Med. **179**:1923.
- Hanissian, S.H. and R.S. Geha (1997) Immunity **6**:379.