

Human PTH1R/PTHR1 Biotinylated Antibody

Antigen Affinity-purified Polyclonal Sheep IgG Catalog Number: BAF5709

DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human PTH1R/PTHR1 in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human PTH1R/PTHR1 Tyr23-Met189 Accession # Q03431
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.
Please Note: Optimal dilute	ions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website. Recommended Sample Concentration
Western Blot	0.1 μg/mL Recombinant Human PTH1R/PTHR1
PREPARATION AND	STORAGE
Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	
	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

BACKGROUND

PTH1 R (Parathyroid hormone receptor # 1; also PTH/PTHr receptor) is a predicted 66-68 kDa member of the G-protein coupled receptor family # 2. It is expressed on a variety of cell types, including hepatocytes, renal epithelium, smooth muscle cells, and osteoblasts plus chondrocytes. PTH1 R is a receptor for both PTH and PTHrP, and PTH binding promotes Ca⁺⁺-release from bone, mediated by osteoclast formation. Mature human PTH1 R is a 7-transmembrane glycoprotein 567 amino acids (aa) in length. It contains a long, ligand-binding N-terminal extracellular region (aa 27-188) and a 130 aa cytoplasmic C-terminal domain. Single aa changes such as Gly121Glu, Ala122Thr and Arg255His can impair PTH1 R signaling. Over aa 1-189, human PTH1 R shares 88% aa identity with mouse PTH1 R.

6 months, -20 to -70 °C under sterile conditions after reconstitution.