

**DESCRIPTION**

<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse OSM R $\beta$ in Western blots.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant mouse OSM R $\beta$ Glu24-Leu738 Accession # O70458
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.1 $\mu$ g/mL	Recombinant Mouse OSM R $\beta$ Fc Chimera (Catalog # 662-OR)

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<p><b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b></p> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

Oncostatin M (OSM) is a member of the IL-6 family of cytokines that share the gp130 as a common signal transducing receptor subunit. Human OSM signals through two types of human OSM receptor complexes: the type I complex comprising the leukemia inhibitory factor receptor beta (LIF R $\beta$ ) and gp130, the type II complex made up of OSM receptor beta (OSM R $\beta$ ) and gp130. In contrast, mouse OSM signals only through the mouse OSM R $\beta$  and gp130 complex. Mouse OSM R $\beta$  cDNA encodes a 971 amino acid (aa) residue type I transmembrane protein which contains a 23 aa residue signal peptide, an extracellular domain of 714 aa, a transmembrane domain of 20 aa and a 214 aa cytoplasmic domain. Mouse OSM R $\beta$  alone binds mOSM with low-affinity, but forms a high-affinity binding complex in the presence of gp130. Mouse OSM R $\beta$  is 55% identical at the amino acid sequence level to human OSM R $\beta$ .

**References:**

1. Lindberg, R.A. *et al.* (1998) *Mol. Cell. Biol.* **18**:3357.
2. Tanaka, M. *et al.* (1999) *Blood* **93**:804.