

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse CD30 Ligand in ELISAs and Western blots. In sandwich immunoassays, less than 0.05% cross-reactivity with recombinant human (rh) CD40 Ligand, rhFas Ligand, rhTNF- α , recombinant mouse TNF- α , recombinant rat TNF- α , and rhTNF- β is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse CD30 Ligand Gln68-Asp239 Accession # P32972
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 μ g/mL	Recombinant Mouse CD30 Ligand/TNFSF8 (Catalog # 732-CL)
Mouse CD30 Ligand/TNFSF8 Sandwich Immunoassay		Reagent
ELISA Capture	2-8 μ g/mL	Mouse CD30 Ligand/TNFSF8 Antibody (Catalog # MAB732)
ELISA Detection	0.1-0.4 μ g/mL	Mouse CD30 Ligand/TNFSF8 Biotinylated Antibody (Catalog # BAF732)
Standard		Recombinant Mouse CD30 Ligand/TNFSF8 (Catalog # 732-CL)

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CD30 Ligand (CD30L)/TNFSF8 is a type II membrane protein belonging to the TNF superfamily. CD30L is expressed on the cell surface of activated T cells, B cells, and monocytes. The protein is also constitutively expressed on granulocytes and medullary thymic epithelial cells. The specific receptor for CD30L is CD30/TNFRSF8, a type I transmembrane glycoprotein belonging to the TNF receptor superfamily. CD30 was originally identified as a cell surface antigen of Hodgkin's and Reed-Sternberg cells using the monoclonal antibody Ki-1. CD30 is also expressed on different non-Hodgkin's lymphomas, virus-infected T and B cells, and on normal T and B cells after activation. Among T cells, CD30 is preferentially expressed on a subset of T cells producing Th2-type cytokines and on CD4⁺/CD8⁺ thymocytes that coexpress CD45RO and IL-4 receptor. CD30 ligation by CD30L mediates pleiotropic effects including cell proliferation, activation, differentiation and cell death by apoptosis. CD30 can act as a co-stimulatory molecule in thymic negative selection and may also play a critical role in the pathophysiology of Hodgkin's disease and other CD30⁺ lymphomas.

References:

1. Brunangelo, F. *et al.* (1995) *Blood* **85**:1.
2. Gruss, H-J. and F. Herrmann (1996) *Leukemia and Lymphoma* **20**:397.
3. Chiarle, R. *et al.* (1999) *J. Immunol.* **163**:194.