



Biotinylated Anti-mouse IL-1 RI Antibody

ORDERING INFORMATION

Catalog Number: BAF771

Lot Number: EXA01

Size: 50 µg

Formulation: 0.2 µm filtered solution in PBS with BSA

Storage: -20° C

Reconstitution: sterile 0.1% BSA in TBS

Specificity: rmlL-1 RI

Immunogen: NS0-derived rmlL-1 RI extracellular domain

Ig Type: goat IgG

Application: Western blot

Preparation

Produced in goats immunized with purified, NS0-derived, recombinant mouse interleukin 1 receptor I (rmlL-1 RI) extracellular domain. Mouse IL-1 RI specific IgG was purified by mouse IL-1 RI affinity chromatography and then biotinylated.

Formulation

Lyophilized from a 0.2 µm filtered solution in phosphate-buffered saline (PBS) containing 50 µg of bovine serum albumin (BSA) per 1 µg of antibody.

Reconstitution

Reconstitute with sterile Tris-buffered saline pH 7.3 (20 mM Trizma base, 150 mM NaCl) containing 0.1% BSA. If 1 mL of buffer is used, the antibody concentration will be 50 µg/mL.

Storage

Lyophilized samples are stable for twelve months from date of receipt when stored at -20° C to -70° C. Upon reconstitution, the antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Reconstituted antibody can also be aliquotted and stored frozen at -20° C to -70° C **in a manual defrost freezer** for six months without detectable loss of activity. **Avoid repeated freeze-thaw cycles.**

Specificity

This antibody has been selected for use as a detection antibody in mouse IL-1 RI western blots.

Application

Western Blot - This antibody can be used at 0.1 - 0.2 µg/mL with the appropriate secondary reagents to detect mouse IL-1 RI. The detection limit for rmlL-1 RI is approximately 5 ng/lane under non-reducing and reducing conditions. In this format, this antibody shows approximately 35% cross-reactivity with rmlL-18 R (IL-1 R5) and rhIL-1 RI and 20% cross-reactivity with rmlL-18 Rβ (IL-1 R7) and rrlL-1 Rrp2 (IL-1 R6) and less than 1% cross-reactivity with rmlL-1 RII.

Optimal dilutions should be determined by each laboratory for each application.