

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human XCR1 in Western blots.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human XCR1 Met1-Thr31, Ser89-Lys103, His168-Val190, Phe251-Tyr267 Accession # P46094
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.1 µg/mL	Recombinant Human XCR1
<b>Flow Cytometry</b>	2.5 µg/10 <sup>6</sup> cells	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

XCR1, also known as GPR5 and lymphotactin/SCM-1 (single cysteine motif 1) receptor, is a 38 kDa member of the G-protein coupled receptor 1 family. It binds XCL1/lymphotactin/SCM-1α and XCL2/SCM-1β. In addition, human herpesvirus 8 (HHV8) encodes two viral chemokines vCCL2/vMIP-II and vCCL3/vMIP-III that function as an antagonist and a highly selective agonist, respectively, for XCR1. XCR1 is expressed on neutrophils, CD8<sup>+</sup> T cells, NK cells, CD4<sup>+</sup> T cells and B cells. Human XCR1 is a 333 amino acid (aa), 7-transmembrane molecule. It contains a 32 aa N-terminus that lacks glycosylation sites and a 42 aa C-terminal cytoplasmic tail. Over the extracellular regions used for immunization, human XCR1 shares 62%, 54% and 64% aa identity with canine, mouse and porcine XCR1, respectively.