

DESCRIPTION

Source	<i>E. coli</i> -derived human Noggin protein Gln28 - Cys232 with an N-terminal Met Accession # Q13253.1
N-terminal Sequence Analysis	Met-Gln28
Structure / Form	Disulfide-linked homodimer
Predicted Molecular Mass	23 kDa (monomer)




SPECIFICATIONS

SDS-PAGE	27 kDa, under reducing conditions.
Activity	Measured by its ability to inhibit BMP-4-induced alkaline phosphatase production by ATDC5 mouse chondrogenic cells. The ED ₅₀ for this effect is 0.0200-0.240 µg/mL
Endotoxin Level	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>97%, by SDS-PAGE with quantitative densitometry by Coomassie® Blue Staining.
Formulation	Lyophilized from a 0.2 µm filtered solution in Sodium Acetate with Trehalose. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 500 µg/mL in water.
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 3 months, -20 to -70 °C under sterile conditions after reconstitution.

DATA



BACKGROUND

Noggin is a secreted homodimeric glycoprotein that is an antagonist of bone morphogenetic proteins (BMPs) (1, 2). Human Noggin cDNA encodes a 232 amino acid (aa) precursor protein; cleavage of a 19 aa signal peptide generates the 213 aa mature protein which contains an N-terminal acidic region, a central basic heparin-binding segment and a C-terminal cysteine-knot structure (2). Secreted Noggin probably remains close to the cell surface due to its binding of heparin-containing proteoglycans (3). Noggin is very highly conserved among vertebrates, such that mature human Noggin shares 99%, 99%, 98%, 97% and 89% aa sequence identity with mouse, rat, bovine, equine and chicken Noggin, respectively. Noggin binds some BMPs such as BMP-4 with high affinity and others such as BMP-7 with lower affinity. It antagonizes BMP bioactivities by blocking epitopes on BMPs that are needed for binding to both type I and type II receptors (2, 4). During embryogenesis, Noggin antagonizes specific BMPs at defined times, for example, during neural tube, somite and cardiomyocyte growth and patterning (5-7). During skeletal development, Noggin prevents chondrocyte hyperplasia, thus allowing proper formation of joints (4). Mutations within the cysteine-knot region of human Noggin are linked to multiple types of skeletal dysplasias that result in apical joint fusions (8). Noggin is expressed in defined areas of the adult central nervous system and peripheral tissues such as lung, skeletal muscle and skin (1). During culture of human embryonic stem cells (hESC) or neural stem cells under certain conditions, addition of Noggin to antagonize BMP activity may allow stem cells to proliferate while maintaining their undifferentiated state, or alternatively, to differentiate into dopaminergic neurons (6, 9 - 13). Noggin also appears to maintain adult stem cell populations in-vivo, for example, maintaining neural stem cells within the hippocampus (13).

References:

1. Valenzuela, D.M. *et al.* (1995) *J. Neurosci.* **15**:6077.
2. Groppe, J. *et al.* (2002) *Nature* **420**:636.
3. Paine-Saunders, S *et al.* (2002) *J. Biol. Chem.* **277**:2089.
4. Brunet, L. J. *et al.* (1998) *Science* **280**:1455.
5. McMahon, J. A. *et al.* (1998) *Genes Dev.* **12**:1438.
6. Itsykson, P. *et al.* (2005) *Mol. Cell. Neurosci.* **30**:24.
7. Yuasa, S. *et al.* (2005) *Nat. Biotechnol.* **23**:607.
8. Gong, Y. *et al.* (1999) *Nat. Genet.* **21**:302.
9. Xu, R.-H. *et al.* (2005) *Nat. Methods* **2**:185.
10. Wang, G. *et al.* (2005) *Biochem. Biophys. Res. Commun.* **330**:934.
11. Chaturvedi, G. *et al.* (2009) *Cell Prolif.* **42**:425.
12. Chiba, S. *et al.* (2008) *Stem Cells* **26**:2810.
13. Bonaguidi, M.A. *et al.* (2008) *J. Neurosci.* **28**:9194.