

**DESCRIPTION**

<b>Source</b>	Chinese Hamster Ovary cell line, CHO-derived	
	Human IL-17A (Gly24-Ala155) Accession # Q16552	
	Human IL-17F (Arg21-GLN163) Accession # Q96PD4	
	N-terminus	C-terminus

**N-terminal Sequence Analysis** Gly24(IL-17A) & Arg21(IL-17F)

**Structure / Form** Disulfide-linked heterodimer, Biotinylated via sugars

**Predicted Molecular Mass** 15 kDa (unlabeled IL-17A), 15 kDa (unlabeled IL-17F)

**SPECIFICATIONS**

**Activity** Measured by its ability to induce IL-6 secretion by NIH-3T3 mouse embryonic fibroblast cells. The ED<sub>50</sub> for this effect is 0.4-2 ng/mL.

**Endotoxin Level** <1.0 EU per 1 µg of the protein by the LAL method.

**Purity** >90%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

**Formulation** Lyophilized from a 0.2 µm filtered solution in HCl with Trehalose. See Certificate of Analysis for details.

**PREPARATION AND STORAGE**


**Reconstitution** Reconstitute at 100 µg/mL in 4 mM HCl.

**Shipping** The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

**Stability & Storage** Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.

**DATA**



**BACKGROUND**

Human IL-17A/F is an approximately 40 kDa, secreted, disulfide-linked heterodimeric glycoprotein comprised of two members of the IL-17 family of cytokines, IL-17A and IL-17F (1, 2). Members of this family demonstrate a structural motif termed a cysteine knot that also characterizes a large superfamily of growth factors. Although most cysteine knot superfamily members use three intrachain disulfide bonds to create a knot, IL-17 family molecules generate the same structural form with only two disulfide links (3-5). Mature human IL-17A and IL-17F share 61% and 56% amino acid sequence identity with mouse IL-17A and IL-17F, respectively. They share 50% aa sequence identity with each other. IL-17A/F and the IL-17A and IL-17F homodimers are produced by IL-23 activated Th17 cells (1, 6-10). The widely expressed receptors IL-17 RA and IL-17 RC form a heterodimer for the binding of IL-17A and IL-17F, as well as the heterodimeric IL-17A/F (6, 11, 12). IL-17A/F is a biologically active protein that induces chemokine production and airway neutrophilia with intermediate potency between IL-17A (most potent) and IL-17F (least potent) (7, 12). It is up-regulated in immune cells during inflammatory arthritis and contributes to disease severity (13).

**References:**

1. Wright, J.F. *et al.* (2007) *J. Biol. Chem.* **282**:13447.
2. Chang, S.H. and C. Dong (2007) *Cell Res.* **17**:435.
3. Kawaguchi, M. *et al.* (2004) *J. Allergy Clin. Immunol.* **114**:1265.
4. Kolls, J.K. and A. Linden (2004) *Immunity* **21**:467.
5. Gu, C. *et al.* (2013) *Cytokine* **64**:477.
6. Cheung, P.F.Y. *et al.* (2008) *J. Immunol.* **180**:5625.
7. Liang, S.C. *et al.* (2007) *J. Immunol.* **179**:7791.
8. Ouyang, W. *et al.* (2008) *Immunity* **28**:454.
9. Steinman, L. (2007) *Nat. Med.* **13**:139.
10. Hunter, C.A. (2005) *Nat. Rev. Immunol.* **5**:521.
11. Kuestner, R.E. *et al.* (2007) *J. Immunol.* **179**:5462.
12. Wright, J.F. *et al.* (2008) *J. Immunol.* **181**:2799.
13. Sarkar, S. *et al.* (2014) *Clin. Exp. Immunol.* **177**:652.