

MATERIAL DATA SHEET

Recombinant Mouse His6 USP18

Cat. # E-604

Ubiquitin-like carboxyl-terminal hydrolase 18 (USP18) is a specialized cysteine protease with a predicted molecular weight of 42 kDa. USP18 from *Mus musculus* is a member of the peptidase C19 family and shares 70% amino acid identity with its human ortholog. USP18 plays an important role in antiviral responses and interferon-mediated signaling pathways. Mice lacking the gene (USP18^{-/-}) are hypersensitive to interferon, have shortened lifespans and pathologic abnormalities of the brain. While it was initially speculated that the cause of these phenotypes stems from the accumulation of ISG15-conjugated intracellular proteins (a target for USP18 catalytic function), it was later demonstrated that USP18 has multiple roles, at least one of which is independent of its ISGylase activity.

Product Information

Quantity:	50 µg
MW:	38 kDa
Source:	<i>Spodoptera frugiperda</i> , Sf21 (baculovirus)-derived Contains an N-terminal 6-His tag, Ser47 - Ser368 Accession # Q9WTV6
Stock:	X mg/ml (X µM) in 50 mM HEPES pH 7.5, 200 mM NaCl, 10% (v/v) Glycerol, 5 mM DTT and 1 mM EDTA
Purity:	>98%, by SDS-PAGE under reducing conditions and visualized by Colloidal Coomassie® Blue stain.

Use & Storage

Use:	Reaction conditions will need to be optimized for each specific application. We recommend an initial recombinant murine USP18 concentration of 100-250 nM when using ISG15-AMC (UL-553) as a substrate.
Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">• 24 months from date of receipt, -70 °C as supplied.• 3 months, -70 °C under sterile conditions after opening.

Literature

References:

1. Goldmann, T. *et al.* (2015) EMBO J. **34**: 1612
2. Ketscher, L. & Knobloch, K.P. (2015) Cytokine **76**: 569
3. Knobloch, K.P. *et al.* (2005) Mol. Cell Biol. **25**: 11030
4. Speer, S.D. *et al.* (2016) Nat. Commun. doi: 10.1038/ncomms11496

For research use only. Not for use in humans.