

Human β -2 Adrenergic R/ADRB2 Alexa Fluor® 350-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 586107

Catalog Number: FAB100401U

100 μ g

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human β -2 Adrenergic R/ADRB2 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 586107
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	NS0 mouse myeloma cell line transfected with human β -2 Adrenergic R/ADRB2 Accession # P07550
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 μ g/10 ⁶ cells	HEK293 Human Cell Line Transfected with Human β -2 Adrenergic R/ADRB2 and eGFP

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Beta-2 adrenoceptor (ADRB2) is a 413 amino acids adrenergic receptor expressed in smooth muscle and metabolic tissues. Beta-adrenergic receptors belong to the seven-transmembrane domain/GPCR family of receptors and mediate the catecholamine-induced activation of adenylate cyclase through the action of G proteins. Activation of ADRB2 induces a decrease in gastrointestinal motility, bronchodilation, vasodilation in skeletal and cardiac muscle, and glycogenolysis in liver. Agonists of ADRB2 are most widely used for the treatment of asthma. In complex with beta-arrestin-1 and c-src, the beta-2 adrenergic receptor activates MAP kinases ERK1(MAPK3) and ERK2 (MAPK1). Expression of the beta-2 adrenoceptor has been reported in adipose, blood, brain, heart, lung, nose, pancreas, skeletal muscle, skin, and vessel.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.