

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse LAIR1 in direct ELISAs.
Source	Monoclonal Rabbit IgG Clone # 2459B
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived mouse LAIR1 Gln22-Tyr141 & Ser25-Tyr141 Accession # Q8BG84
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Blockade of Receptor-ligand Interaction Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Leukocyte-associated Ig-like receptor-1 (LAIR1; also CD305) is a 46 kDa inhibitory receptor of the Ig superfamily that is structurally related to inhibitory members of KIR and ILT/CD85 families (1-3). It is expressed on immune cells, including NK cells, T cells, B cells, monocytes, immature neutrophils, dendritic cells and most thymocytes (2-4). The 253 amino acid (aa) type I transmembrane (TM) protein contains a 21 aa signal sequence, a 124 aa extracellular domain (ECD), a 20 aa TM domain and a 98 aa cytoplasmic domain. The ECD includes one C2-type Ig-like domain and two potential N-linked glycosylation sites. Tyrosine phosphorylation of two cytoplasmic ITIM motifs results in recruitment of phosphatases and down-regulation of signaling through activating receptors (2, 3, 5). Crosslinking of LAIR1 inhibits processes such as B cell receptor-mediated activation, NK cell and T cell cytotoxicity and basophil degranulation (1-3). Four mouse LAIR1 splice variants have been identified, but it is not known whether they are expressed as proteins (3). LAIR1b, which is the major alternate transcript, lacks most of the ECD. Of the minor transcripts, LAIR1c lacks a signal sequence, and LAIR1d and 1e lack a TM sequence. All mouse splice forms are identical in the last 90 aa of the cytoplasmic domain. LAIR1 shows high-affinity binding of collagens that results in inhibition of degranulation in a basophilic leukemia cell line (6). Human and mouse LAIR1 ECD share only 32% aa identity but, where studied, sites of expression and activity are similar (3-6). Mouse LAIR1 ECD also shares 62%, 31% and 28% aa identity with rat, canine, and bovine orthologs, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.